

REDUCIBLE AND TOROIDAL DEHN FILLINGS WITH DISTANCE 3

SUNGMO KANG

ABSTRACT. If a simple 3-manifold M admits a reducible and a toroidal Dehn filling, the distance between the filling slopes is known to be bounded by three. In this paper, we classify all manifolds which admit a reducible Dehn filling and a toroidal Dehn filling with distance 3.

Keywords: Dehn filling; Reducible; Toroidal; Tangle filling.

1. INTRODUCTION

Let M be a compact connected orientable 3-manifold with a torus boundary component $\partial_0 M$ and r a *slope*, the isotopy class of an essential simple closed curve, on $\partial_0 M$. The manifold obtained by r -Dehn filling is defined to be $M(r) = M \cup V$, where V is a solid torus glued to M along $\partial_0 M$ so that r bounds a disk in V .

We say that M is *simple* if it contains no essential sphere, torus, disk or annulus. For a pair of slopes r_1 and r_2 on $\partial_0 M$, the distance $\Delta(r_1, r_2)$ denotes their minimal geometric intersection number. For a simple manifold M , if both $M(r_1)$ and $M(r_2)$ fail to be simple, then the upper bounds for $\Delta(r_1, r_2)$ have been established in various cases. In particular, for the case of reducible and toroidal Dehn fillings, Oh [15] and independently Wu [17] showed that for a simple manifold M , if $M(r_1)$ is reducible and $M(r_2)$ is toroidal then $\Delta(r_1, r_2) \leq 3$. Eudave-Muñoz and Wu [2] have explicitly described an infinite family of manifolds M realizing $\Delta(r_1, r_2) = 3$ in terms of tangle arguments. More specifically, M is the double branched cover of one of the tangles illustrated in Figure 1 where p is an integer no less than 3. However, p can be extended to any integer no less than 2.

The goal of this paper is to show that the manifolds M described above are the only examples of simple manifolds admitting a reducible Dehn filling and a toroidal Dehn filling with distance 3. The main results of this paper are the following.

Theorem 1.1. *Let M be a simple 3-manifold with a torus boundary component $\partial_0 M$ such that $M(r_1)$ is reducible, $M(r_2)$ is toroidal and*




FIGURE 1. The tangles and the $1/0$ - and $1/3$ -rational tangle fillings used in Theorem 1.1.

$\Delta(r_1, r_2) = 3$. Then M is the double branched cover of one of the tangles with $p \geq 2$ as described in Figure 1 (i). $M(r_1)$ corresponding to the $1/3$ -rational tangle filling is a connected sum of two lens spaces $L(3, 1)$ and $L(2, 1)$, and $M(r_2)$ corresponding to the $1/0$ -rational tangle filling is toroidal and is not Seifert fibered. See Figure 1 (ii), (iii).

In order to prove Theorem 1.1, we need the following.

Theorem 1.2. *Let M be a simple 3-manifold as in Theorem 1.1. Then $M(r_2)$ is the union of M_1 and M_2 , where M_i , $i = 1, 2$, is a Seifert fibered space over a disk with two exceptional fibers, along an incompressible torus. Furthermore, the fibers of M_1 and M_2 intersect exactly once on the incompressible torus i.e. $M(r_2)$ is not Seifert fibered.*

In [12], Lee showed that if $M(r_1)$ is reducible and $M(r_2)$ contains a Klein bottle, then M is uniquely determined. That is, M is homeomorphic to $W(6)$, which is obtained from the exterior W of the Whitehead link by performing Dehn filling on one of its boundary components with slope 6 under the standard meridian-longitude coordinates. Let $M = W(6)$. Then it is well-known that $M(1)$ is reducible, and $M(4)$ is

toroidal and contains a Klein bottle. Moreover, using the fact that W is strongly invertible, we can easily see that M is the double branched cover of the tangle illustrated in Figure 1 (i) with $p = 2$, say T , and $M(1)$ and $M(4)$ are the double branched covers of T with the $1/3$ - and $1/0$ -rational tangles filled respectively. Notice that the toroidal Dehn filling $M(4)$ is the union of M_1 and M_2 , where M_i , $i = 1, 2$, is a Seifert fibered space over a disk with two exceptional fibers, along an incompressible torus and the fibers of M_1 and M_2 intersect exactly once on the incompressible torus i.e. $M(4)$ is not Seifert fibered. We have shown that if $M(r_2)$ contains a Klein bottle, then Theorem 1.1 and Theorem 1.2 hold.

Therefore, throughout this paper we may assume that $M(r_2)$ does not contain a Klein bottle. Then Theorem 1.1 and Theorem 1.2 follow immediately from Theorem 7.1 and Theorem 5.17 respectively.

As an immediate consequence of Theorem 1.2, we have the following.

Corollary 1.3. *Let M be a simple manifold with a torus boundary component. If $M(r_1)$ is reducible and $M(r_2)$ is a toroidal Seifert fibered manifold, then $\Delta(r_1, r_2) \leq 2$.*

It is unknown whether or not the upper bound 2 in Corollary 1.3 is the best possible.

2. THE INTERSECTION GRAPHS

From now on we assume that M is a simple 3-manifold with a torus boundary component $\partial_0 M$ and that r_1 and r_2 are slopes on $\partial_0 M$ such that $M(r_1)$ is reducible and $M(r_2)$ contains an essential torus and $\Delta(r_1, r_2) = 3$.

Over all reducing spheres in $M(r_1)$ which intersect the attached solid torus V_1 in a family of meridian disks, we choose a 2-sphere \widehat{F}_1 so that $F_1 = \widehat{F}_1 \cap M$ has the minimal number, say n_1 , of boundary components. Similarly let \widehat{F}_2 be an essential torus in $M(r_2)$ which intersects the attached solid torus V_2 in a family of meridian disks, the number of which, say n_2 , is minimal over all such surfaces and let $F_2 = \widehat{F}_2 \cap M$. Let u_1, u_2, \dots, u_{n_1} be the disks of $\widehat{F}_1 \cap V_1$, labeled as they appear along V_1 . Similarly let v_1, v_2, \dots, v_{n_2} be the disks of $\widehat{F}_2 \cap V_2$. Then F_1 is an essential planar surface, and F_2 is an essential punctured torus in M . We may assume that F_1 and F_2 intersect transversely and the number of components of $F_1 \cap F_2$ is minimal over all such surfaces. Then no circle component of $F_1 \cap F_2$ bounds a disk in either F_1 or F_2 and no arc component is boundary-parallel in either F_1 or F_2 . In this section, the subscripts i and j will denote 1 or 2 with the convention that

if they both appear, then $\{i, j\} = \{1, 2\}$. The components of ∂F_i are numbered $1, 2, \dots, n_i$ according to the labels of the corresponding disks of $\widehat{F}_i \cap V_i$. We obtain a graph G_i in \widehat{F}_i by taking as the (fat) vertices of G_i the disks in $\widehat{F}_i \cap V_i$ and as the edges of G_i the arc components of $F_1 \cap F_2$ in F_i . Each endpoint of an edge of G_i has a label, that is, the label of the corresponding component of ∂F_j , $i \neq j$. Since each component of ∂F_i intersects each component of ∂F_j in Δ points, the labels $1, 2, \dots, n_j$ appear in order around each vertex of G_i repeatedly Δ times.

For a graph G , the *reduced graph* \overline{G} of G is defined to be the graph obtained from G by amalgamating each family of mutually parallel edges into a single edge. For an edge α of \overline{G} , the *weight* of α , denoted by $w(\alpha)$, is the number of edges of G in α .

Orient all components of ∂F_i so that they are mutually homologous on $\partial_0 M$, $i = 1, 2$. Let e be an edge in G_i . Since e is a properly embedded arc in F_i , it has a disk neighborhood D in F_i with $\partial D = a \cup b \cup c \cup d$, where a and c are arcs in ∂F_i with induced orientation from ∂F_i . On D , if ∂D can be oriented so that a and c have the same orientation as that induced from ∂F_i , then e is called *positive*, otherwise *negative*. See Figure 2. Then we have the parity rule.

An edge is positive on one graph if and only if it is negative on the other graph.




FIGURE 2. Orientations on an edge.

Orient the core of V_i . Then we can give a sign to each vertex of G_i according to the sign of its intersection with the core of V_i . Two vertices (possibly equal) of G_i are called *parallel* if they have the same sign, otherwise *antiparallel*. A positive edge connects parallel vertices while a negative one connects antiparallel vertices. Let G_i^+ denote the subgraph of G_i consisting of all the vertices and all the positive edges of G_i .

Let G be G_1 or G_2 and let x be a label of G . An x -edge is an edge of G with the label x at one endpoint. An x -cycle is a cycle of positive x -edges which can be oriented so that the tail of each edge has the label x . A cycle in G is a *Scharlemann cycle* if it bounds a disk face, and the edges in the cycle are all positive and have the same label pair. If the label pair is $\{x, y\}$, then we refer to such a Scharlemann cycle as an (x, y) -Scharlemann cycle. In particular, a Scharlemann cycle of length 2 is called an S -cycle.

Lemma 2.1. *If G_i has a Scharlemann cycle, then \widehat{F}_j is separating.*

Proof. This follows from [17, Lemma 1.2]. \square

3. BINARY FACES AND THE SUBGRAPH Λ OF G_1^+

In this section, we will define binary faces in G_1^+ and see some properties of binary faces. Since the arguments here are exactly the same as in [10] and in [12] at the same time, we use the terminologies and several Lemmas without proof described in [10] and in [12]. We may assume that $M(r_2)$ is irreducible by [8].

Lemma 3.1. *G_2 has exactly two vertices and the two vertices of G_2 are antiparallel.*

Proof. By Theorem 1.3 in [14], G_2 has exactly two vertices v_1 and v_2 . Assume that v_1 and v_2 are parallel. Then all the edges of G_2 are positive. Since M is simple, $n_1 \geq 3$. Thus by Lemma 1.5 in [17] we can take a label x which is not a label of a Scharlemann cycle in G_2 . Consider the subgraph Γ of G_2 consisting of all vertices and all x -edges of G_2 . Let V, E and F be the numbers of vertices, edges and disk faces of Γ , respectively. Since $V < E$, we have $0 = \chi(\widehat{F}_2) \leq V - E + F < F$, so Γ contains a disk face, which is an x -face in G_2 . This contradicts [14, Theorem 4.4]. \square

We assume without loss of generality that the ordering of the labels around v_1 is anticlockwise while the ordering around v_2 is clockwise. Since the two vertices of G_2 are antiparallel, an Euler characteristic argument on G_1^+ shows that there is a disk face in G_1^+ , which is a Scharlemann cycle. Then \widehat{F}_2 is separating in $M(r_2)$ by Lemma 2.1. We will call one side of \widehat{F}_2 in $M(r_2)$ the *W(hite)* side, denoted by \widehat{M}_W and the other the *B(lack)* side, denoted by \widehat{M}_B . Then F_2 has a W side and a B side in M , which are denoted by M_W, M_B respectively. The faces of G_1 are called W and B according to which side of F_2 a neighborhood of their boundary lies.

G_2 has at most 6 *edge classes*, the isotopy classes in \widehat{F}_2 relative to the fat vertices of G_2 by [3, Lemma 5.2]. We label these edge classes $\alpha, \beta, \gamma, \delta, \epsilon, \epsilon'$ as in Figure 3. An edge in G_1 or G_2 is called an η -*edge* if, being regarded as an edge in G_2 , it lies in class η , $\eta \in \{\alpha, \beta, \gamma, \delta, \epsilon, \epsilon'\}$. The ϵ -edges and ϵ' -edges are positive in G_2 while the others are negative by Lemma 3.1. Since v_1 and v_2 have the same valency, $w(\epsilon) = w(\epsilon')$. In Figure 3, an ϵ -edge has two endpoints on ∂v_1 . If the endpoint lies between the edge class β and the edge class γ , we say that the endpoint lies in class ϵ_N , otherwise ϵ_S . If the endpoint of an ϵ' -edge lies between the edge class α and the edge class γ , we say that the endpoint lies in class ϵ'_N , otherwise ϵ'_S .




FIGURE 3. 6 edge classes on \widehat{F}_2 .

Two of those (non-loop) edge classes are said to be *adjacent* if ignoring the edge classes ϵ, ϵ' , the corresponding edges are adjacent in \overline{G}_2 . They are *non-adjacent* otherwise. Thus, as in Figure 3, the pair $\{\alpha, \gamma\}$ and the pair $\{\beta, \delta\}$ are non-adjacent. Let λ, μ be two elements of $\{\alpha, \beta, \gamma, \delta\}$. Then a (λ, μ) *face* of G_1 is one whose edges lie in classes λ, μ on G_2 . Lemma 3.1 of [7] shows that any such face must have edges in both classes λ and μ . We call any such face a *binary face* of G_1 .

A (λ, μ) face f is λ -*good* if no two consecutive edges in the boundary of f belong to class λ . A (λ, μ) face is *good* if it is either λ -good or μ -good. A *bad* face is a binary face that is not good. To indicate the coloring of an (λ, μ) face we will often call it an $X(\lambda, \mu)$ face where X is to take the value B or W .

If c is an X corner of G_1 for which the edge incident to label 1 is in class λ and the edge incident to label 2 is in class μ , we call c an $X(\lambda, \mu)$

corner, where $X \in \{B, W\}$. Then notice that a bad $X(\lambda, \mu)$ face of G_1 must contain $X(\lambda, \lambda)$, $X(\lambda, \mu)$, $X(\mu, \lambda)$ and $X(\mu, \mu)$ corners. For negative edges, there are four different types of corners i.e. $X(\epsilon_N, \epsilon'_N)$, $X(\epsilon_N, \epsilon'_S)$, $X(\epsilon_S, \epsilon'_N)$ and $X(\epsilon_S, \epsilon'_S)$ corners.

Then we have the theorem which plays an important role in this paper.

Theorem 3.2. *Assume that G_1 contains a good $X(\lambda, \mu)$ face of length n . Then the X side of \widehat{F}_2 in $M(r_2)$ is a Seifert fibered space over the disk with exactly two exceptional fibers, one of which has order n . Moreover the Seifert fiber of this fibration is the curve on \widehat{F}_2 formed by $\lambda \cup \mu$.*

Proof. This is exactly Theorem 4.1 in [10]. □

The following lemmas are shown in Section 3 of [10].

Lemma 3.3. *Let c_1, c_2, \dots, c_n be a collection of X corners of G_1 , $X \in \{B, W\}$. The anticlockwise ordering of the endpoints of the edges of these corners on vertex 1 of G_2 is the same as the clockwise ordering of the edge endpoints on vertex 2 of G_2 .*

Lemma 3.4. *G_1 cannot contain a vertex v with two edges in the same edge class incident to v at the same label.*

Lemma 3.5. *Given $X \in \{B, W\}$, G_1 does not contain $X(\lambda, \lambda)$ corners for three distinct edge classes λ .*

Lemma 3.6. *Suppose that G_1 contains a bad $X(\alpha, \delta)$ face. Then there exists $\lambda \in \{\alpha, \delta\}$ such that, for any edge e of G_1 in class β or γ , the edges of G_1 adjacent to e on the X -side are in class λ . More generally, the analogous statement holds whenever $\{\alpha, \delta\}$ is replaced by any pair of adjacent edge classes.*

Lemma 3.7. *G_1 does not contain bad faces of the same color on distinct edge class pairs.*

Lemma 3.8. *If G_1 contains a good $X(\lambda, \mu)$ face, then any binary X -face of G_1 on an edge class pair distinct but not disjoint from $\{\lambda, \mu\}$ is bad.*

[12, Proposition 4.3] says that G_1^+ contains a connected subgraph Λ satisfying the following properties;

- (1) for all vertices u_x of Λ but at most one (an exceptional vertex), there are an edge with edge class α or β , and an edge with edge class γ or δ in Λ which are incident to u_x with label i for each $i = 1, 2$;

- (2) for the exceptional vertex u_{x_0} , if it exists, there are at least two edges in Λ incident to u_{x_0} ;
- (3) there is a disk $D_\Lambda \subset \widehat{F}_1$ such that $\text{Int} D_\Lambda \cap G_1^+ = \Lambda$; and
- (4) Λ has no cut vertex.

A vertex of Λ is a *boundary vertex* if there is an arc connecting it to ∂D_Λ whose interior is disjoint from Λ , and an *interior vertex* otherwise. Similarly for a *boundary edge* and an *interior edge* of Λ .

Lemma 3.9. *Λ contains a bigon.*

Proof. This follows from the proof of [12, Lemma 4.8]. \square

Lemma 3.10. *Λ contains a face bounded only by α - or β -edges, or only by γ - or δ -edges. In other words, Λ contains a $X(\lambda, \mu)$ face where $\{\lambda, \mu\} = \{\alpha, \beta\}$ or $\{\gamma, \delta\}$ and $X \in \{B, W\}$.*

Proof. This follows from [12, Lemma 4.6]. \square



By Lemma 3.10, from now on, throughout the paper we may assume without loss of generality that Λ contains a $B(\alpha, \beta)$ face.

Let \widehat{H}_B (resp. \widehat{H}_W) be the intersection of V_2 and \widehat{M}_B (resp. \widehat{M}_W). Let H_B (resp. H_W) be the intersection of V_2 and M_B (resp. M_W). Then $V_2 = \widehat{H}_B \cup \widehat{H}_W$ and $\partial V_2 = H_B \cup H_W$. For the purpose of Sections 4 and 5 there are two equivalent figures of H_B or H_W as in Figure 4 where corners of faces in G_1^+ appear. Figure 4 (ii) is obtained by cutting H_B or H_W along the arc a in Figure 4 (i) such that one endpoint of a lies between the edge class α and the edge class ϵ_S at ∂v_1 and the other lies between the edge class α and the edge class β at ∂v_2 .

Lemma 3.11. *A $B(\alpha, \beta)$ face is good.*

Proof. Suppose a $B(\alpha, \beta)$ face is bad. Then it contains $B(\alpha, \alpha)$, $B(\alpha, \beta)$, $B(\beta, \alpha)$, $B(\beta, \beta)$ corners. These corners lie on H_B as in Figure 5. There are two cases. First, we assume Figure 5 (i). Figure 5 (ii) is similar. It follows from Figure 5 (i) that $\omega(\epsilon') + \omega(\delta) + \omega(\gamma) + \omega(\epsilon') \leq \omega(\beta)$. On the other hand, $\omega(\alpha) + \omega(\epsilon') + \omega(\delta) + \omega(\gamma) + \omega(\epsilon') + \omega(\beta) = 3n_1$ since the number of all the labels around a vertex in G_2 is $3n_1$. Therefore, combining this equation with the above inequality, we get the inequality $\omega(\alpha) + 2\omega(\beta) \geq 3n_1$. This implies that either $\omega(\alpha) \geq n_1$ or $\omega(\beta) \geq n_1$, which contradicts [17, Lemma 1.5]. \square

Lemma 3.12. *The black side \widehat{M}_B of $M(r_2)$ is a Seifert fibered space over the disk with exactly two exceptional fibers. Moreover the Seifert fiber of this fibration is the curve on \widehat{F}_2 formed by $\alpha \cup \beta$.*

FIGURE 4. Two equivalent figures of H_W or H_B .FIGURE 5. The intersection of H_B and the corners of a $B(\alpha, \beta)$ face.

Proof. Λ contains a $B(\alpha, \beta)$ face, which we denote by f . Let λ_i , $i = 1, \dots, n$, be the corner of f and $\partial\lambda_i^j$ the endpoint of λ_i at the label j , $j = 1, 2$. Let $I(v_j)$ be the shortest interval on ∂v_j containing all $\partial\lambda_i^j$ in its interior. Relabel i so that $\partial I(v_1) = \{\partial\lambda_1^1, \partial\lambda_n^1\}$. It follows from [3, Lemma 2.4] that $\partial I(v_2) = \{\partial\lambda_1^2, \partial\lambda_n^2\}$. Since f is good by Lemma 3.11, there exists a disk D in H_B bounded by four arcs a_1, a_2, b_1, b_2 where

a_i is the arc of ∂v_i containing $I(v_i)$, $i = 1, 2$, and b_j is the parallel arc to λ_j in H_B , $j = 1, 2$, so that D contains all the corners of f .

We can take an annulus A on F_2 containing all the edges of f so that $A \cup D$ is an once punctured torus on M_B and the two boundary components of A are homotopic to the curve on \widehat{F}_2 formed by $\alpha \cup \beta$. See Figure 3. Since ∂f is non-separating on $A \cup D$, we can get a disk D' by surgering $A \cup D$ along f . Push D' slightly into the interior of M_B so that D' is a properly embedded disk in M_B . Notice that $\partial D'$ contains the two arcs b_1, b_2 . An orientation of $\partial D'$ induces orientations of b_1, b_2 that are opposite on H_B . Hence shrinking \widehat{H}_B to its core yields a properly embedded annulus A' in the black side \widehat{M}_B of \widehat{F}_2 . In addition, A' separates \widehat{M}_B into two manifolds M_1 and M_2 , both of which have a torus boundary i.e. $\widehat{M}_B = M_1 \cup_{A'} M_2$. We can observe that the core of \widehat{H}_B lies on A' . Therefore by isotoping the core of V_2 slightly off ∂M_1 and ∂M_2 , we can show that M_1 and M_2 are solid tori by the choice of \widehat{F}_2 . Since \widehat{F}_2 is essential in $M(r_2)$, the black side \widehat{M}_B is a Seifert fibered space over the disk with two exceptional fibers and the fiber is the core of A' and hence the core of A , as desired. \square

Remark: In the proof of Lemma 3.12, the core of \widehat{H}_B , that is, the black part of V_2 , lies on the annulus which separates the Seifert fibered space \widehat{M}_B into two solid tori. We will use this fact to prove Lemma 3.13.

Before proving the next lemma, we observe that the edge classes of the three edges of a trigon in Λ cannot be all distinct by Lemma 3.3 and cannot be all same by Lemma 3.1 of [7].

Lemma 3.13. *If Λ contains a white bigon or a white trigon, the two edge classes of the edges of a white bigon or a white trigon are λ and μ where $\{\lambda, \mu\} = \{\alpha, \delta\}, \{\alpha, \gamma\}, \{\beta, \delta\}$ or $\{\beta, \gamma\}$.*

Proof. Let f be a white bigon in Λ . f is necessarily a good $W(\lambda, \mu)$ face. Suppose $\{\lambda, \mu\} = \{\alpha, \beta\}$ or $\{\gamma, \delta\}$. Then applying the same argument as in the proof of Lemma 3.12 using f , we see that the white side \widehat{M}_W is a Seifert fibered space over the disk with exactly two exceptional fibers and the Seifert fiber of this fibration is the curve on \widehat{F}_2 formed by $\lambda \cup \mu$. In other words, $\widehat{M}_W = M_3 \cup_A M_4$ where M_3 and M_4 are solid tori and A is a properly embedded annulus in \widehat{M}_W , and the core of A is the fiber. On the other hand, the black side $\widehat{M}_B = M_1 \cup_{A'} M_2$ where M_1 and M_2 are solid tori and the core of A' is the fiber. By the *Remark* below Lemma 3.12, the core of V_2 lies on $A \cup A'$. Furthermore since two fibers of \widehat{M}_B and \widehat{M}_W coincide on \widehat{F}_2 , we can isotope A and

A' on \widehat{F}_2 fixing the core of V_2 so that $A \cup A'$ is a torus \widehat{T} , say. Note that either side of \widehat{T} is a Seifert fibered space over the disk with two exceptional fibers, which implies that \widehat{T} is essential in $M(r_2)$. Pushing the core of V_2 off \widehat{T} , we get an essential torus in $M(r_2)$ which misses the core of V_2 , which is a contradiction to the choice of \widehat{F}_2 . Therefore $\{\lambda, \mu\} = \{\alpha, \delta\}, \{\alpha, \gamma\}, \{\beta, \delta\}$ or $\{\beta, \gamma\}$.

Let g be a white trigon in Λ . Hence g is also necessarily a good $W(\lambda, \mu)$ face. Applying the same argument as in the case of a bigon above, we complete the proof. \square

Lemma 3.14. *Any two black bigons in G_1^+ have the same pair of edge classes. Furthermore the pair is either $\{\alpha, \beta\}$ or $\{\gamma, \delta\}$.*

Proof. Lemma 3.3 implies that no two black bigons of G_1^+ have distinct and non-disjoint pairs of edge classes, nor have the pair $\{\{\alpha, \gamma\}, \{\beta, \delta\}\}$. Therefore the only possible different pairs of edge classes of two bigons are $\{\{\alpha, \beta\}, \{\gamma, \delta\}\}, \{\{\alpha, \delta\}, \{\beta, \gamma\}\}$. In either case, we construct two Möbius bands from the two bigons by shrinking \widehat{H}_B to its core. Furthermore, the two boundaries of the two Möbius bands cobound an annulus in \widehat{F}_2 , which gives a Klein bottle in $M(r_2)$, a contradiction to the assumption that $M(r_2)$ does not contain a Klein bottle. Hence any two black bigons in G_1^+ have the same pair of edge classes.

Now we show that the pair is either $\{\alpha, \beta\}$ or $\{\gamma, \delta\}$. If there is a black bigon in G_1^+ with the edge class pair $\{\lambda, \mu\}$, then the black side \widehat{M}_B of \widehat{F}_2 in $M(r_2)$ is a Seifert fibered space over the disk with exactly two exceptional fibers and the Seifert fiber of this fibration is the curve on \widehat{F}_2 formed by $\lambda \cup \mu$ by Theorem 3.2. However the fiber of \widehat{M}_B is the curve on \widehat{F}_2 formed by $\alpha \cup \beta$ by Lemma 3.12. Since $M(r_2)$ does not contain a Klein bottle, the Seifert fibration of \widehat{M}_B is unique by [11, Theorem VI.18]. Therefore the edge class pair $\{\lambda, \mu\}$ of the bigon must be either $\{\alpha, \beta\}$ or $\{\gamma, \delta\}$. \square

The next theorem is the main result of the combinatorial arguments in this paper. To prove this, we use Lemma 3.17 and the main theorems in Sections 4 and 5.

Theorem 3.15. *Λ contains a $W(\lambda, \mu)$ bigon, where $\{\lambda, \mu\} = \{\alpha, \delta\}, \{\alpha, \gamma\}, \{\beta, \delta\}$ or $\{\beta, \gamma\}$.*

Proof. Lemma 3.9 says that Λ contains a bigon f . If f is white, then we are done by Lemma 3.13.

Suppose Λ does not contain a white bigon. Then f is black. By Lemma 3.14, we may assume that f is a $B(\alpha, \beta)$ bigon. Lemma 3.17

says that Λ contains a trigon. On the other hand, Theorem 4.1 and Theorem 5.1 imply that Λ does not contain a trigon. This contradiction shows that Λ contains a white bigon as desired. \square

To prove Lemma 3.17, Theorem 4.1 and Theorem 5.1, we assume that Λ contains a $B(\alpha, \beta)$ bigon and does not contain a white bigon.

Lemma 3.14 enables us to apply Lemmas 4.10 and 4.11 of [12] without change.

Lemma 3.16. *All interior vertices of Λ have valency at least 4 in $\overline{\Lambda}$ and all boundary vertices of Λ but the exceptional vertex have valency at least 3 in $\overline{\Lambda}$.*

Proof. This is Lemma 4.10 of [12]. \square

Lemma 3.17. *Λ contains a trigon.*

Proof. This is Lemma 4.11 of [12]. \square

Lemma 3.18. *Λ contains an interior vertex.*

Proof. Suppose that Λ has no interior vertices. Lemma 3.16 guarantees that the number of vertices of Λ is greater than 3. Put Λ on an abstract disk D so that Λ lies in the interior of D and insert an edge connecting each vertex of Λ to ∂D . By taking the reduced graph of the resulting graph, we obtain a graph Γ satisfying the supposition of [1, Lemma 2.6.5]. In addition, since the number of vertices of Γ is greater than 3, Γ must satisfy (*) in the proof of [1, Lemma 2.6.5]. In other words, Γ contains two vertices of valency at most 3. However this is impossible since all vertices of Γ , possibly except one, have valency at least 4 by Lemma 3.16. \square

In Section 4, we will show that Λ does not contain a black trigon and in Section 5, we will show that Λ does not contain a white trigon.

4. A $B(\alpha, \beta)$ BIGON AND A BLACK TRIGON

Throughout this section, we assume that Λ contains a $B(\alpha, \beta)$ bigon and does not contain a white bigon. Our goal in this section is to show the following.

Theorem 4.1. *If Λ contains a $B(\alpha, \beta)$ bigon and does not contain a white bigon, then Λ does not contain a black trigon.*

The main idea is to define two dual graphs of Λ and use the fact that there is no triple of non-isomorphic disk faces in Λ and apply the index equation $\sum_{\text{vertices}} I(v) + \sum_{\text{faces}} I(f) = 2$.

To prove Theorem 4.1, we suppose throughout this section that Λ contains a black trigon.

Lemma 4.2. *All black trigons of G_1^+ have the same edge classes, i.e. two γ -edges and one δ -edge, say.*

Proof. Let g be a black trigon of G_1^+ . Lemma 3.3 excludes the case that the three edges of g have distinct edge classes. Thus g has two λ -edges and one μ -edge. Since g is a good $B(\lambda, \mu)$ face, we have the Seifert fibration of the black side \widehat{M}_B resulting from g whose fiber is represented by $\{\lambda, \mu\}$. Since $M(r_2)$ does not contain a Klein bottle, by the uniqueness of the Seifert fibration, $\{\lambda, \mu\} = \{\alpha, \beta\}$ or $\{\gamma, \delta\}$ by Lemma 3.12.

Since Λ contains a $B(\alpha, \beta)$ bigon, there are two corners, a $B(\alpha, \beta)$ corner and a $B(\beta, \alpha)$ corner. Then Lemma 3.3 eliminates the case that $\{\lambda, \mu\} = \{\alpha, \beta\}$. However Lemma 3.3 also excludes the case that Λ contains simultaneously two trigons with two γ -edges and one δ -edge, and two δ -edges and one γ -edge respectively. Hence we may assume that all trigons of G_1^+ have the same edge classes, i.e. without loss of generality two γ -edges and one δ -edge. \square

Lemma 4.3. *Λ cannot contain a $B(\lambda, \mu)$ corner, where $(\lambda, \mu) = (\alpha, \delta)$, (β, γ) , (γ, β) , (δ, α) , or (δ, δ) .*

Proof. Λ contains a black bigon f_1 with one α -edge and one β -edge, and contains a black trigon f_2 with two γ -edges and one δ -edge. Consider the intersection of all the corners of f_1 and f_2 , and H_B . See Figure 6. Then the above black corners are impossible. \square

Since Λ contains a black face and a white face simultaneously by Lemma 3.16, M_B and M_W are handlebodies of genus 2 from [7, Lemma 8.3].

Traveling around the boundary of a disk face of G_1^+ gives rise to a cyclic sequence of edge classes. We shall say that two disk faces of G_1^+ of the same color are *isomorphic* if the cyclic sequences obtained by traveling in some directions are equal. See [9, Section 5].

Recall that $H_B = V_2 \cap M_B$ and $H_W = V_2 \cap M_W$. Then $\partial M_B = H_B \cup F_2$ and $\partial M_W = H_W \cup F_2$. Since all the vertices of Λ are parallel, each face of Λ is a non-separating disk in M_B or M_W . Note that any two faces of the same color in Λ are disjoint.

Lemma 4.4. *If two disk faces of G_1^+ are parallel in M_B or M_W , then they are isomorphic.*

Proof. This is Lemma 4.13 of [12]. \square

Theorem 4.5. *There is no triple of mutually non-isomorphic black disk faces in G_1^+ , and there is no triple of mutually non-isomorphic white disk faces in G_1^+ .*




FIGURE 6. A $B(\alpha, \beta)$ bigon, a $B(\gamma, \delta)$ trigon and the intersection on H_B .

To prove this theorem, we will show that there is a white binary disk face by using dual graphs of Λ .

For the purpose of the proof of Theorem 4.5, we assume for the time being that $G_1 = G_1^+$. Then every vertex of Λ except an exceptional vertex has valency 6 since all edges of G_1 are positive.

Now we define a dual graph Λ^* to Λ as in [10, Section 3]. To do this, it is convenient to regard Λ as a graph in S^2 , rather than a disk. In other words, we add an additional *outside face* to the set of faces of Λ . Then a *vertex* of Λ^* is a point in the interior of each face of Λ and a point of the outside face. An *edge* of Λ^* is the edge connecting the vertices of Λ^* corresponding to the faces of Λ on either side of each edge of Λ , and meeting each edge transversely in a single point.

We give two orientations ω and ω' to the edges of Λ^* as follows;

ω : an edge of Λ^* dual to an edge of Λ in an edge class $\in \{\alpha, \delta\}$ (resp. $\{\beta, \gamma\}$) is oriented from the *W-side* to the *B-side* (resp. from the *B-side* to the *W-side*).

ω' : an edge of Λ^* dual to an edge of Λ in an edge class $\in \{\beta, \delta\}$ (resp. $\{\alpha, \gamma\}$) is oriented from the *W-side* to the *B-side* (resp. from the *B-side* to the *W-side*).

See Figure 7.

Note. For an edge e of Λ in the boundary of the outside face, we regard the side of e contained in the outside face as being locally colored with the color opposite to that of the face of Λ that contains e .




FIGURE 7. Two orientations ω and ω' to the edges of Λ^* .

As defined in [6], the *index* of a vertex v and a face f of a graph in S^2 with oriented edges is defined by $I(v) = 1 - s(v)/2$ and $I(f) = \chi(f) - s(f)/2$ where $s(v)$ and $s(f)$ are switches at v and of f respectively. A *sink* or a *source* is a vertex of index 1, and a *cycle* is a face of index 1. We say an *exceptional cycle* if it is dual to an exceptional vertex.

Lemma 4.6. Λ^* does not contain a cycle except an exceptional cycle with respect to ω, ω' .

Proof. Suppose v is a non-exceptional vertex dual to a cycle with respect to ω . Since v has valency 6, the edge classes of three edges which are incident to v at the label 1 belong to either $\{\alpha, \delta\}$ or $\{\beta, \gamma\}$. Therefore two of them have the same edge class at the same label. This contradicts Lemma 3.4. The same argument applies to ω' . \square

Let s and s' be the number of sinks and sources of Λ^* at vertices dual to the faces of Λ (i.e. we exclude the outside face) with respect to ω, ω' respectively.

Lemma 4.7. $s, s' > 0$.

Proof. By Lemma 4.6, the only possible cycle is an exceptional cycle. However if Λ^* has an exceptional cycle, then it cannot contain the sink

or source dual to the outside face at the same time. Hence it follows from the equation $\sum_{\text{vertices}} I(v) + \sum_{\text{faces}} I(f) = 2$ that $s, s' > 0$. \square

A sink or source of ω, ω' in Λ^* that does not correspond to the outside face of Λ is a binary face of Λ .

Lemma 4.8. *There is a white binary face in Λ dual to a sink or source with respect to ω .*

Proof. By Lemma 4.7, there is a binary face in Λ dual to a sink or source with respect to ω . We show that this binary face is white.

Suppose that a binary face in Λ dual to a sink or source with respect to ω is black. Then such a black binary face is either a $B(\alpha, \delta)$ face or a $B(\beta, \gamma)$ face. The former contains a $B(\alpha, \delta)$ corner and the latter contains a $B(\beta, \gamma)$ corner. This contradicts Lemma 4.3. This completes the proof. \square

So far, we have shown that under the assumption that $G_1 = G_1^+$ Lemma 4.8 holds. Now we are ready to prove Theorem 4.5.

Proof of Theorem 4.5. We follow the proof of Lemma 4.16 of [12]. Let f_1, f_2, f_3 be three mutually non-isomorphic black faces of G_1^+ . Then these faces cut M_B into two 3-balls by Lemma 4.4.

Let f be a black face of G_1 other than f_1, f_2, f_3 . Since f lies in the complement of $f_1 \cup f_2 \cup f_3$ in M_B , f must be a disk, otherwise it would be compressible in M_B , so F_1 would be compressible in M , a contradiction to the fact that F_1 is incompressible. Each component of $\partial f \cap F_2$ is an edge of G_2 . ∂f must be an essential curve in $\partial M_B = F_2 \cup H_B$, otherwise some component of $\partial f \cap F_2$ would be a trivial loop in G_2 . Hence f is an essential disk in M_B , so it must be parallel to one of the faces f_1, f_2 and f_3 . Therefore any black face of G_1 is a disk face isomorphic to one of f_1, f_2, f_3 by Lemma 4.4. It follows that all the edges of G_1 are positive, i.e. $G_1 = G_1^+$.

Next, we will show that any white face of G_1 is a disk face. Note that if every white face of G_1 is a disk face, then G_1 is connected. Suppose a white face g of G_1 is not a disk face. Lemma 4.8 guarantees the existence of a white binary disk face g_1 . Furthermore, there must be a white disk face g_2 other than g_1 in Λ which is non-isomorphic to g_1 , otherwise there are three edges in the same edge class incident to an interior vertex, contradicting Lemma 3.4. Thus the complement of $g_1 \cup g_2$ in M_W is a 3-ball. So, g is compressible in M_W , which implies that F_1 is compressible, a contradiction.

We have shown that $G_1 = G_1^+$ and G_1 is connected. Therefore, F_1 is nonseparating, which contradicts [13, Theorem 1.1]. This completes

the proof that there is no triple of mutually non-isomorphic black disk faces in G_1^+ . Similarly, we can apply the same argument to show that there is no triple of non-isomorphic white disk faces, by using a $B(\alpha, \beta)$ bigon. \square

Now we go back to the subgraph Λ as in Section 3 and continue with the proof of Theorem 4.1. Since Λ contains a $B(\alpha, \beta)$ bigon and a $B(\gamma, \delta)$ trigon with two γ -edges and one δ -edge by Lemma 4.2, Theorem 4.5 implies the following.

Lemma 4.9. *Any black disk face of G_1^+ is isomorphic to either a $B(\alpha, \beta)$ bigon or a $B(\gamma, \delta)$ trigon with two γ -edges and one δ -edge.*

We define the dual graph Λ^* with two orientations ω, ω' as in the case $G_1 = G_1^+$. The cycle dual to an interior vertex in Λ is said to be an *ordinary cycle*. The cycle dual to a boundary vertex (except an exceptional vertex) is called a *boundary cycle*. The cycle dual to an exceptional vertex is said to be an *exceptional cycle*. Thus when we say a boundary cycle dual to a boundary vertex, we do not allow it to be the cycle dual to an exceptional vertex.

Lemma 4.10. Λ^* *does not contain a sink or source dual to a black face of Λ with respect to ω, ω' .*

Proof. Lemma 4.9 says that any black disk face is either a $B(\alpha, \beta)$ bigon or a $B(\gamma, \delta)$ trigon with two γ -edges and one δ -edge. However, either black disk face is not dual to a sink or source with respect to ω, ω' . \square

It is guaranteed by Lemma 3.18 that Λ contains an interior vertex. Note that there are only four distinct types of interior vertices. See Figure 8.




FIGURE 8. Four distinct types of interior vertices.

Lemma 4.11. Λ^* does not contain an ordinary cycle with respect to ω, ω' .

Proof. Since any interior vertex has valency 6, this follows from the proof of Lemma 4.6. \square

Lemma 4.12. A boundary cycle dual to a boundary vertex with respect to ω, ω' is one of those illustrated in Figure 9 and Figure 10 respectively.

Proof. If a boundary vertex v dual to a boundary cycle has valency 5 or 6, then at least two edges incident to v have the same edge class at the same label, contradicting Lemma 3.4. Hence any boundary vertex dual to a boundary cycle has valency 4. Furthermore, any boundary vertex has two α - or β -edges and also two γ - or δ -edges by the property (1) of Λ . Then the proof follows from Lemma 4.9. \square





FIGURE 9. Boundary cycles with respect to ω .


Lemma 4.13. Λ^* does not contain a boundary cycle with respect to ω' .

Proof. Assume for contradiction that Λ contains a boundary cycle dual to a boundary vertex u_x . In G_2 , the label x appears either three times around v_1 at the ends of α -, γ - and ϵ -edges and three times around v_2 at the ends of β -, δ - and ϵ' -edges, or three times around v_1 at the ends of β -, δ - and ϵ -edges and three times around v_2 at the ends of α -, γ - and ϵ' -edges. The odd-numbered types in Figure 10 have the first case and the even-numbered types have the second case.

Claim. In G_2 , the label x appears at ϵ_N (resp. ϵ_S) of the ϵ -edge class at v_1 if and only if it appears at ϵ'_S (resp. ϵ'_N) of the ϵ' -edge class at v_2 .

FIGURE 10. Boundary cycles with respect to ω' .

Proof. Assume the first case i.e. the label x appears three times around the vertex v_1 at the ends of α -, γ - and ϵ -edges and three times around the vertex v_2 at the ends of β -, δ - and ϵ' -edges. The second case is similar.

FIGURE 11. Three occurrences of label x on v_1 and v_2 .

Suppose, for example, that x appears at ϵ_N and at ϵ'_N of ϵ - and ϵ' -edges respectively at the same time. See Figure 11. At the vertex v_1 the edge classes δ and ϵ don't have the label x . Then the sum of the weights of δ and ϵ is less than n_1 . On the other hand, at the vertex v_2 the edge classes δ and ϵ' have the label x twice, which implies that the sum of the weights of δ and ϵ' is greater than n_1 . However, these two inequalities conflict since the weights of ϵ and ϵ' are same. \square

Assume that the boundary vertex u_x is dual to one of the types $(i) \sim (iv)$. Then we have a $B(\epsilon, \epsilon')$ corner. In Figure 6 of Lemma 4.3, (ϵ, ϵ') should be $(\epsilon_N, \epsilon'_N)$ or $(\epsilon_S, \epsilon'_S)$. This contradicts the Claim.

For the type (v) , we have two black corners, a $B(\gamma, \epsilon')$ corner, a $B(\epsilon, \delta)$ corner. However, from Figure 6 of Lemma 4.3, (ϵ, ϵ') should be $(\epsilon_N, \epsilon'_N)$, which is a contradiction to the Claim. The other types are also impossible by the same argument. This completes the proof. \square

Note. The Claim in the proof of Lemma 4.13 doesn't work with the orientation ω .

Lemma 4.14. Λ^* does not contain a boundary cycle with respect to ω .

Proof. Lemmas 4.10, 4.11, 4.13 and the equation $\sum_{\text{vertices}} I(v) + \sum_{\text{faces}} I(f) = 2$ imply that there is a sink or source dual to a white disk face with respect to ω' . In other words, there is a $W(\beta, \delta)$ or $W(\alpha, \gamma)$ face. There are eight types of boundary cycles with respect to ω as in Figure 9. The first four types have a $B(\epsilon, \epsilon')$ corner. However, by Figure 6, (ϵ, ϵ') must be either $(\epsilon_N, \epsilon'_N)$ or $(\epsilon_S, \epsilon'_S)$. Assume we have the type (i) . If we let u_x be a boundary vertex of the type (i) , then the label x appears three times around the vertex v_1 at the ends of β -, γ - and ϵ -edges and three times around the vertex v_2 at the ends of α -, δ - and ϵ' -edge. Applying a similar argument as in the Claim in Lemma 4.13, (ϵ, ϵ') must be $(\epsilon_S, \epsilon'_S)$. Thus, we have a $W(\beta, \delta)$ corner, a $W(\epsilon_S, \alpha)$ corner, and a $W(\gamma, \epsilon'_S)$ corner. Since we have a $W(\beta, \delta)$ or $W(\alpha, \gamma)$ face, there are $W(\delta, \delta)$ or $W(\beta, \beta)$ corners in a $W(\beta, \delta)$ face by the Remark below the proof of Lemma 4.4 in [10], and there are $W(\alpha, \gamma)$ corners in a $W(\alpha, \gamma)$ face. Then neither a $W(\beta, \beta)$ corner, a $W(\delta, \delta)$ corner nor a $W(\alpha, \gamma)$ corner satisfies Lemma 3.3 with $W(\beta, \delta)$, $W(\epsilon_S, \alpha)$, and $W(\gamma, \epsilon'_S)$ corners. Therefore type (i) can't happen. A similar argument applies to the other types (ii) , (iii) and (iv) .

For the type (v) , we have a $W(\beta, \delta)$ corner, a $W(\gamma, \alpha)$ corner and a $W(\epsilon, \epsilon')$ corner. Then Lemma 3.3 forces (ϵ, ϵ') to be either $(\epsilon_N, \epsilon'_S)$ or $(\epsilon_N, \epsilon'_S)$. If $(\epsilon, \epsilon') = (\epsilon_N, \epsilon'_S)$, the boundary vertex has a $B(\gamma, \epsilon'_S)$ corner, which is impossible by Figure 6. If $(\epsilon, \epsilon') = (\epsilon_S, \epsilon'_N)$, then the boundary vertex has a $B(\epsilon_S, \delta)$ corner, which is also impossible by Figure 6. For the other types, we can apply a similar argument. \square

Proof of Theorem 4.1. Lemmas 4.10, 4.11 and 4.13 guarantee that Λ contains a $W(\lambda_1, \lambda_2)$ face where $\{\lambda_1, \lambda_2\} = \{\beta, \delta\}$ or $\{\alpha, \gamma\}$ by the index equation $\sum_{\text{vertices}} I(v) + \sum_{\text{faces}} I(f) = 2$. Also Lemmas 4.10, 4.11 and 4.14 guarantee that Λ contains a $W(\mu_1, \mu_2)$ face where $\{\mu_1, \mu_2\} = \{\alpha, \delta\}$ or $\{\beta, \gamma\}$. Hence a $W(\lambda_1, \lambda_2)$ face and a $W(\mu_1, \mu_2)$ face are the only two non-isomorphic white disk faces in Λ by Theorem 4.5.

If $\{\lambda_1, \lambda_2\} = \{\beta, \delta\}$ and $\{\mu_1, \mu_2\} = \{\alpha, \delta\}$, then no γ -edges are incident to interior vertices, which is impossible since there are only four types of interior vertices. See Figure 8. Similarly for the case that $\{\lambda_1, \lambda_2\} = \{\beta, \delta\}$ and $\{\mu_1, \mu_2\} = \{\beta, \gamma\}$, in which case no α -edges are incident to interior vertices, and the case that $\{\lambda_1, \lambda_2\} = \{\alpha, \gamma\}$ and $\{\mu_1, \mu_2\} = \{\alpha, \delta\}$, in which case no β -edges are incident to interior vertices.

Then the only remaining case is that $\{\lambda_1, \lambda_2\} = \{\alpha, \gamma\}$ and $\{\mu_1, \mu_2\} = \{\beta, \gamma\}$, i.e. Λ has a $W(\alpha, \gamma)$ face and a $W(\beta, \gamma)$ face, in which case no δ -edges are incident to interior vertices, which is possible when an interior vertex is of type (i) or (ii). These faces cannot be both bad by Lemma 3.5 and cannot be both good by Lemma 3.8.




FIGURE 12. The intersection of the white corners of a bad $W(\beta, \gamma)$ face and H_W , and the local configuration at an interior vertex of type (i).

Assume that a $W(\alpha, \gamma)$ face is good and a $W(\beta, \gamma)$ face is bad. Then every interior vertex is of type (i). Consider the intersection of the white corners of a bad $W(\beta, \gamma)$ face and H_W where there are two possibilities as in Figure 12 (i), (ii), and the local configuration at an interior vertex of type (i) as in Figure 12 (iii). However, Figure 12 (ii) is impossible since there is a $W(\alpha, \gamma)$ corner. In Figure 12 (iii) e must be γ from Figure 12 (i). Then the vertex v has two γ -edges at the same label, which is a contradiction to Lemma 3.4.

Assume that a $W(\alpha, \gamma)$ face is bad and a $W(\beta, \gamma)$ face is good. Then every interior vertex is of type (ii). Consider the intersection of the white corners of a $W(\alpha, \gamma)$ face and H_W and the local configuration at an interior vertex of type (ii). See Figure 13. There are two possibilities for the intersection of the white corners of a $W(\alpha, \gamma)$ face and H_W as in Figure 13 (i), (ii). Figure 13 (ii) is impossible since there is a $W(\beta, \gamma)$

corner. In Figure 13 (iii) suppose that v_1 is not an exceptional vertex. Then e_1 must be an α -edge and must be a boundary edge since the δ -edge adjacent to e_1 is a boundary edge. This implies that e_2 is an interior edge and then a β -edge. Also, e_3 must be a γ -edge from Figure 13 (i). This is a contradiction to Lemma 3.4. If v_1 is an exceptional vertex, consider the vertex v_2 in Figure 13 (iii) and apply the same argument. Hence we have finished the proof of Theorem 4.1. \square




FIGURE 13. The intersection of the white corners of a $W(\alpha, \gamma)$ face and H_W , and the local configuration at an interior vertex of type (ii).

5. A $B(\alpha, \beta)$ BIGON AND A WHITE TRIGON.

In this section, we assume that Λ contains a $B(\alpha, \beta)$ bigon and does not contain a white bigon. Then the main goal of this section is to prove the following theorem.

Theorem 5.1. *If Λ contains a $B(\alpha, \beta)$ bigon and does not contain a white bigon, then Λ does not contain a white trigon.*

Suppose that Λ contains a white trigon. Then Lemma 3.13 says that a white trigon is either a $W(\alpha, \delta)$, $W(\beta, \gamma)$, $W(\beta, \delta)$ or $W(\alpha, \gamma)$ trigon.

Note that Theorem 4.5 still holds by using a white trigon instead of a white binary disk face in the proof of Theorem 4.5. Furthermore, by Lemma 3.18, there is an interior vertex in Λ .

First, assume that there is a $W(\alpha, \delta)$ trigon with two δ -edges and one α -edge in Λ .

Lemma 5.2. *Λ^* does not contain a boundary cycle with respect to ω' .*

Proof. Consider the intersection of the white corners of a $W(\alpha, \delta)$ trigon and H_W where there are two cases as in Figure 14 (i), (ii), and the intersection of the black corners of a $B(\alpha, \beta)$ bigon and H_B as in Figure 14 (iii). Let v be a boundary vertex dual to a boundary cycle with respect to ω' . There are two types of local configuration at v as shown in Figure 15 (i), (ii), where e_1, e_2, e_3 and e_4 are positive edges and e_5 and e_6 are negative edges.




FIGURE 14.




FIGURE 15.

Assume we have Figure 14 (ii). Then the only possible white corner with one β -edge is a $W(\beta, \delta)$ corner or a $W(\delta, \beta)$ corner, which makes a boundary vertex not to be a cycle with respect to ω' since a β -edge and a δ -edge have the same orientation with respect to ω' . Now assume Figure 14 (i).

1) The first type as in Figure 15 (i). Note that $(e_5, e_6) = (\epsilon_N, \epsilon'_S)$ from the intersection of the white corners of a $W(\alpha, \delta)$ trigon and H_W as shown in Figure 14 (i).

Assume there is a clockwise boundary cycle. There are only two cases for e_1 i.e. e_1 is either α or γ . First if $e_1 = \alpha$, then e_2 must be δ from Figure 14 (i) since there is a $W(\epsilon_N, \epsilon'_S)$ corner. Then $e_3 = \gamma$ and $e_4 = \beta$. Thus, we have a $B(\gamma, \delta)$ corner and a $B(\epsilon_N, \beta)$ corner. However this cannot happen from the intersection of the black corners of a $B(\alpha, \beta)$ bigon and H_B as in Figure 14 (iii).

Second, if $e_1 = \gamma$, e_2 must be β since there is a $W(\epsilon_N, \epsilon'_S)$ corner. Then $e_3 = \alpha$ and $e_4 = \delta$. Hence we have two black corners, i.e. a $B(\gamma, \epsilon'_S)$ corner and a $B(\epsilon_N, \delta)$ corner. We can observe from Figure 14 (iii) with a $B(\gamma, \epsilon'_S)$ corner and a $B(\epsilon_N, \delta)$ corner inserted that the only positive edge from which a black corner runs to a γ -edge is a γ -edge. Thus if a γ -edge is an interior edge, then there is a face all edges of which are γ -edges, which contradicts Lemma 3.1 of [7]. Consequently, no γ -edge is incident to interior vertices. For an interior vertex u , a δ -edge must be incident to u , otherwise there are three α or β -edges incident to u , which is a contradiction to Lemma 3.4. However at whatever label a δ -edge is incident to u , there is another δ -edge incident to u at the same label from Figure 14 (i) with a $W(\epsilon_N, \epsilon'_S)$ corner inserted and Figure 14 (iii) with a $B(\gamma, \epsilon'_S)$ corner and a $B(\epsilon_N, \delta)$ corner inserted, which is a contradiction to Lemma 3.4. See Figure 15 (iii), (iv).

Assume there is an anticlockwise boundary cycle. If $e_1 = \beta$, then $e_2 = \gamma$, $e_3 = \delta$ and $e_4 = \alpha$ since there is a $W(\epsilon_N, \epsilon'_S)$ corner. Thus, we have a $B(\delta, \gamma)$ corner and a $B(\beta, \epsilon'_S)$ corner. However these two black corners cannot happen at the same time from the intersection of the black corners of a $B(\alpha, \beta)$ bigon and H_B as in Figure 14 (iii). If $e_1 = \delta$, then $e_2 = \alpha$, $e_3 = \beta$ and $e_4 = \gamma$. Hence we have two black corners, i.e. a $B(\epsilon_N, \gamma)$ corner and a $B(\delta, \epsilon'_S)$ corner at v . We can observe that the only positive edge class to which a black corner runs from a γ -edge is a γ -edge. The argument is then completely the same as that of the case of a clockwise boundary cycle.

2) The second type as in Figure 15 (ii). We use a similar argument as in the proof of Lemma 4.13. If we let u_x be a boundary vertex dual to a boundary cycle, then the label x appears either three times around v_1 at the ends of α -, γ - and ϵ -edges and three times around v_2 at the ends of β -, δ - and ϵ' -edges, or three times around v_1 at the ends of β -, δ - and ϵ -edges and three times around v_2 at the ends of α -, γ - and ϵ' -edges. By the Claim in Lemma 4.13, (e_5, e_6) is either $(\epsilon_N, \epsilon'_S)$ or $(\epsilon_S, \epsilon'_N)$. However, one of e_1 and e_4 must be either α or γ . Thus by Figure 14 (i), $(e_5, e_6) = (\epsilon_N, \epsilon'_S)$. Figure 14 (iii) with a $B(\epsilon_N, \epsilon'_S)$ corner inserted shows that if one edge of a black corner is a δ -edge, then the other of the corner must be a β -edge. Therefore there is no boundary cycle with respect to ω' . This completes the proof. \square




FIGURE 16.




FIGURE 17.

Lemma 5.3. *There is no source or sink dual to a white face in Λ^* with respect to ω' .*

Proof. Suppose there is a source or sink dual to a white face with respect to ω' , i.e. there is a $W(\lambda, \mu)$ face where $\{\lambda, \mu\} = \{\alpha, \gamma\}$ or $\{\beta, \delta\}$. Since Λ contains a $W(\alpha, \delta)$ trigon, which is good, Lemma 3.8 implies that a $W(\lambda, \mu)$ face is bad. If $\{\lambda, \mu\} = \{\alpha, \gamma\}$ i.e. Λ contains a bad $W(\alpha, \gamma)$ face, then we have a $W(\alpha, \alpha)$ corner, a $W(\gamma, \gamma)$ corner and a $W(\delta, \delta)$ corner. This is impossible by Lemma 3.5. Therefore we assume that Λ has a bad $W(\beta, \delta)$ face. Note that Theorem 4.5 implies that every white face in Λ is either a $W(\alpha, \delta)$ trigon or a bad $W(\beta, \delta)$ face. Consider the intersection of the white corners of a $W(\alpha, \delta)$ trigon,

the white corners of a bad $W(\beta, \delta)$ face and H_W where there is only one case, and the intersection of the black corners of a $B(\alpha, \beta)$ bigon and H_B . See Figure 16. Since Λ has a $W(\alpha, \delta)$ trigon and a bad $W(\beta, \delta)$ face, there are at least two vertices which have a $W(\delta, \delta)$ corner. Let v be such a non-exceptional vertex.

Claim. *There is only one local configuration at v as shown in Figure 17 (i).*

Proof. If at least five positive edges are incident to v , then there are at least two white corners at v which are corners of white disk faces in Λ . One is a $W(\delta, \delta)$ corner. The other corner must be a $W(\beta, \beta)$ corner because every white face in Λ is either a $W(\alpha, \delta)$ trigon or a bad $W(\beta, \delta)$ face and no vertex has three edges incident to it in the same edge class. The one remaining white corner at v must be a $W(\alpha/\gamma/\epsilon, \alpha/\gamma/\epsilon')$ corner, which is impossible by Figure 16 (i). If four edges incident to v are positive, the local configuration at v must be as in Figure 17 (i) since a $W(\epsilon, \epsilon')$ corner cannot occur by Figure 16 (i). \square

Suppose a $W(\delta, \delta)$ corner of a bad $W(\beta, \delta)$ face doesn't appear at an exceptional vertex. The bad $W(\beta, \delta)$ face contains a vertex u as shown in Figure 17 (ii). Since there is only one local configuration at the vertex containing a $W(\delta, \delta)$ corner, e is positive. However, e cannot be α, γ and δ from Figure 16 (ii) because there is a $B(\epsilon_N, \epsilon'_S)$ corner by the Claim. Also, e cannot be β by Lemma 3.4.

Finally, every $W(\delta, \delta)$ corner of a $W(\beta, \delta)$ face must appear at an exceptional vertex. Therefore there is only one $W(\beta, \delta)$ face X , say. Thus there is only one possible sink or source dual to a white face. Note that an exceptional vertex is not dual to a cycle in Λ^* because it contains a $W(\delta, \delta)$ corner. On the other hand, every white face in Λ is either a $W(\alpha, \delta)$ trigon or a bad $W(\beta, \delta)$ face. Thus there is a unique type of interior vertices as in Figure 17 (iii) since a $B(\epsilon_N, \epsilon'_S)$ corner coming from the Claim and Figure 16 (ii) exclude a $B(\delta, \delta)$ corner. Since there are only two non-isomorphic black disk faces in Λ , one black disk face has an α -, β - and δ -edges coming from the local picture at an interior vertex as shown in Figure 17 (iii). The other black disk face is a $B(\alpha, \beta)$ bigon. Hence there is no sink or source dual to a black disk face. Since there is no boundary cycle by Lemma 5.2, the only possible positive index takes place at X or the outside face. However, the index of an interior vertex is -1 . This is a contradiction to the equation $\sum_{vertices} I(v) + \sum_{faces} I(f) = 2$. \square

Lemma 5.4. *There is no source or sink dual to a black face in Λ^* with respect to ω' .*




FIGURE 18.




FIGURE 19.

Proof. Suppose there is a source or sink dual to a black face with respect to ω' , i.e. there is a $B(\lambda, \mu)$ face where $\{\lambda, \mu\} = \{\alpha, \gamma\}$ or $\{\beta, \delta\}$. Then a $B(\lambda, \mu)$ face must be bad by Lemma 3.8 because of the existence of a $B(\alpha, \beta)$ bigon.

Suppose we have a bad $B(\beta, \delta)$ face. Consider the intersection of the black corners of a $B(\alpha, \beta)$ bigon, the black corners of a bad $B(\beta, \delta)$ face and H_B as shown in Figure 18 (iii), and the intersection of the white corners of a $W(\alpha, \delta)$ trigon and H_W where there are two cases as shown in Figure 18 (i), (ii). By Theorem 4.5, every black face is either a $B(\alpha, \beta)$ bigon or a bad $B(\beta, \delta)$ face. Also, since three edges in the same edge class cannot be incident to a vertex, there are only two types of local configuration at interior vertices. See Figure 19 (i) (ii). Note that Figure 18 (i) excludes the first type of interior vertices because a $W(\alpha, \alpha)$ corner and a $W(\delta, \beta)$ corner cannot exist simultaneously in

Figure 18 (i), and Figure 18 (ii) rules out the second type of interior vertices since no $W(\beta, \beta)$ corner survives in Figure 18 (ii).

Let v be a vertex containing a $B(\beta, \beta)$ corner of a bad $B(\beta, \delta)$ face. Suppose v is a non-exceptional vertex. Now apply a similar argument as in the proof of the Claim in Lemma 5.3 by using Figure 18 (iii). Then the local configuration at v looks like Figure 19 (iii). However, a $W(\epsilon_N, \epsilon'_S)$ corner is impossible because in the case of Figure 18 (i) the second type of interior vertices contains a $W(\beta, \beta)$ corner, and in the case of Figure 18 (ii) a $W(\epsilon_N, \epsilon'_S)$ corner cannot exist.

Consequently, v must be an exceptional vertex and then there is only one bad $B(\beta, \delta)$ face Y , say, containing only one $B(\beta, \beta)$ corner. Note that since v contains a $B(\beta, \beta)$ corner, the exceptional vertex v is not dual to a cycle in Λ^* . We already showed that there are no $W(\beta, \delta)$ faces and no $W(\alpha, \gamma)$ faces in Λ . Furthermore, no $B(\alpha, \gamma)$ faces exist since we have a bad $B(\beta, \delta)$ face. Therefore, there are only two possible sinks or sources in Λ^* . One is dual to Y and the other the outside face.

Since there is no boundary cycle in Λ^* by Lemma 5.2, in order to get a contradiction to the equation $\sum_{vertices} I(v) + \sum_{faces} I(f) = 2$, it suffices to show that either there is a vertex of index -1 or the outside face is not dual to a sink or source. Let u be a vertex containing a $B(\beta, \delta)$ corner of Y . Then u is not an exceptional vertex. Also u is a boundary vertex since the local picture at interior vertices looks like Figure 19 (i) (ii). We can observe that two negative edges are not incident to u since neither a $W(\epsilon, \epsilon')$ corner nor a $B(\epsilon, \epsilon')$ corner exists by Figure 18 with white corners of an interior vertex added. If at most one negative edge is incident to u , it is easy to see using Figure 18 that either u has index -1 or the outside face is not dual to a sink or a source.

Suppose we have a bad $B(\alpha, \gamma)$ face in Λ . Apply a similar argument as in the case of a $B(\beta, \delta)$ face. First use a $B(\alpha, \alpha)$ corner instead of a $B(\beta, \beta)$ corner to show that there is only one $B(\alpha, \gamma)$ face in Λ . However unlike the case of a $B(\beta, \delta)$ face, there is only one type of local configuration at interior vertices and furthermore the index of an interior vertex is -1 . Use the equation $\sum_{vertices} I(v) + \sum_{faces} I(f) = 2$ to get a contradiction. \square

By Lemmas 5.2, 5.3, and 5.4 and the index equation $\sum_{vertices} I(v) + \sum_{faces} I(f) = 2$, we have proved Theorem 5.1 for the case that Λ contains a $W(\alpha, \delta)$ trigon with two δ -edges and one α -edge.

Next, suppose Λ contains a $W(\alpha, \delta)$ trigon with two α -edges and one δ -edge.

Lemma 5.5. Λ^* does not contain a boundary cycle with respect to ω' .

Proof. The proof of this lemma is exactly analogous as that of Lemma 5.2. \square

Lemma 5.6. *There is no source or sink dual to a white face in Λ^* with respect to ω' .*




FIGURE 20.




FIGURE 21.

Proof. Suppose there is a source or sink dual to a white face with respect to ω' , i.e. there is a $W(\lambda, \mu)$ face where $\{\lambda, \mu\} = \{\alpha, \gamma\}$ or $\{\beta, \delta\}$. Then a $W(\lambda, \mu)$ face must be a bad $W(\alpha, \gamma)$ face by Lemma 3.8 and Lemma 3.5. Consider the intersection of the white corners of a $W(\alpha, \delta)$ trigon, the white corners of a bad $W(\alpha, \gamma)$ face and H_W where there is only one case, and the intersection of the black corners of a $B(\alpha, \beta)$ bigon and H_B . See Figure 20. Since Λ contains a $W(\alpha, \delta)$ trigon and a bad $W(\alpha, \gamma)$ face, there are at least two vertices which have a $W(\alpha, \alpha)$ corner. Let v be such a non-exceptional vertex.

Claim. *There is only one local configuration at v as shown in Figure 21 (i).*

Proof. Follow the proof of the Claim in Lemma 5.3 with β, δ replaced by γ, α respectively. \square

Let u be an interior vertex in Λ . Since every white face in Λ is either a $W(\alpha, \delta)$ trigon or a bad $W(\alpha, \gamma)$ face by Theorem 4.5, there are only two possible types of the local configuration at u by Lemma 3.4. See Figure 21 (ii), (iii). However either type is also impossible since a $B(\delta, \gamma)$ corner in Figure 21 (ii) and a $B(\delta, \delta)$ corner in Figure 21 (iii) cannot happen by the existence of a $B(\epsilon_N, \epsilon'_S)$ corner coming from the local configuration at v in the Claim. This completes the proof. \square

Lemma 5.7. *There is no source or sink dual to a black face in Λ^* with respect to ω' .*

Proof. Suppose there is a source or sink dual to a black face with respect to ω' , i.e. there is a $B(\lambda, \mu)$ face where $\{\lambda, \mu\} = \{\alpha, \gamma\}$ or $\{\beta, \delta\}$. Then a $B(\lambda, \mu)$ face must be bad by Lemma 3.8 since Λ contains a $B(\alpha, \beta)$ bigon.




FIGURE 22.

Suppose we have a bad $B(\beta, \delta)$ face. Consider the intersection of the black corners of a $B(\alpha, \beta)$ bigon, the black corners of a bad $B(\beta, \delta)$ face and H_B as in Figure 22 (iii), and the intersection of the white corners of a $W(\alpha, \delta)$ trigon and H_W where there are two cases as in Figure 22 (i), (ii). By Theorem 4.5, every black face is either a $B(\alpha, \beta)$ bigon or a bad $B(\beta, \delta)$ face. Also, since three edges in the same edge class cannot be incident to a vertex, there are only two types of the local configuration at interior vertices. See Figure 23 (i), (ii). Note that the white corners of an interior vertex rule out Figure 22 (ii).




FIGURE 23.

Claim. A bad $B(\beta, \delta)$ face whose $B(\beta, \beta)$ corner appears at a non-exceptional vertex contributes 0 to the sum $\sum_{\text{vertices}} I(v) + \sum_{\text{faces}} I(f)$.

Proof. Let v be a vertex containing a $B(\beta, \beta)$ corner of a bad $B(\beta, \delta)$ face. A similar argument as in the proof of the Claim in Lemma 5.3 shows that the local configuration at v looks like Figure 23 (iii). Then v has index -1 . This completes the proof. \square

Suppose no $B(\beta, \beta)$ corners appear at an exceptional vertex. Lemmas 5.5, 5.6 and the Claim imply that the only possible positive index comes from the cycle dual to an exceptional vertex or the sink or source dual to the outside face. However both cannot happen simultaneously so we get a contradiction to the equation $\sum_{\text{vertices}} I(v) + \sum_{\text{faces}} I(f) = 2$. Hence there is a $B(\beta, \beta)$ corner of a $B(\beta, \delta)$ face which must appear at an exceptional vertex. Then the rest of the proof is pretty similar to the fourth and the fifth paragraphs of the proof of Lemma 5.4.

For the case that there is a bad $B(\alpha, \gamma)$ face in Λ , apply a similar argument as in the case of a $B(\beta, \delta)$ face. \square

By Lemmas 5.5, 5.6, and 5.7 and the index equation $\sum_{\text{vertices}} I(v) + \sum_{\text{faces}} I(f) = 2$, we have proved Theorem 5.1 for the case that Λ contains a $W(\alpha, \delta)$ with two α -edges and one δ -edge.

For the case of a $W(\beta, \gamma)$ trigon, we can apply the same argument as in the case of a $W(\alpha, \delta)$ trigon by interchanging α and β , and by interchanging γ and δ .

Now suppose Λ contains a $W(\beta, \delta)$ trigon with two δ -edges and one β -edge. Here, instead of using the orientation ω' , we will use the orientation ω .

Lemma 5.8. Λ^* does not contain a boundary cycle with respect to ω .




FIGURE 24.




FIGURE 25.

Proof. Consider the intersection of the white corners of a $W(\beta, \delta)$ trigon and H_W , where there are two cases as in Figure 24 (i), (ii), and the intersection of the black corners of a $B(\alpha, \beta)$ bigon and H_B as in Figure 24 (iii). Let v be a boundary vertex dual to a boundary cycle with respect to ω . There are two types of the local configuration at v as shown in Figure 25 where e_1, e_2, e_3 and e_4 are positive edges, and e_5 and e_6 are negative edges.

First we assume Figure 24 (i); then if one edge of a white corner is an α -edge, then the other edge of the corner must be a δ -edge. Since an α -edge and a δ -edge have the same orientation with respect to ω , there is no boundary cycle, as desired.

Second we assume Figure 24 (ii); then

1) The first type as in Figure 25 (i); Note that $(e_5, e_6) = (\epsilon_S, \epsilon'_N)$ from the intersection of the white corners of a $W(\beta, \delta)$ trigon and H_W as in Figure 24 (ii). That is, there is a $W(\epsilon_S, \epsilon'_N)$ corner. Then Figure

24 (ii) with this corner inserted shows that if one edge of a white corner in Λ is an α -edge, then the other edge must be a δ -edge. Therefore, there is no boundary cycle.

2) The second type as in Figure 25 (ii); Assume there is a clockwise boundary cycle. If $e_2 = \gamma$ and $e_4 = \beta$, then $e_1 = \alpha$ and $e_3 = \delta$. In other words, we have a $B(\gamma, \alpha)$ corner and a $B(\beta, \delta)$ corner, which is impossible by Figure 24 (iii). If $e_2 = \beta$ and $e_4 = \gamma$, then e_5 must be δ by Figure 24 (ii), which is absurd because e_5 is a negative edge.

Assume there is an anticlockwise boundary cycle. If $e_1 = \gamma$, e_6 must be δ by Figure 24 (ii), which is absurd. If $e_1 = \beta$ and $e_3 = \gamma$, then $e_2 = \delta$ and $e_4 = \alpha$. In other words, we have a $B(\alpha, \gamma)$ corner and a $B(\delta, \beta)$ corner, which is impossible by Figure 24 (iii). \square

Lemma 5.9. *There is no source or sink dual to a white face in Λ^* with respect to ω .*

Proof. Apply exactly the same argument as that of Lemma 5.3 by interchanging α and β . \square

Lemma 5.10. *There is no source or sink dual to a black face in Λ^* with respect to ω .*

Proof. The proof is exactly the same as that of Lemma 5.4 with α and β interchanged. \square

In conclusion, by Lemmas 5.8, 5.9, and 5.10 and the index equation $\sum_{vertices} I(v) + \sum_{faces} I(f) = 2$, we have proved Theorem 5.1 for the case that Λ contains a $W(\beta, \delta)$ trigon with two δ -edges and one β -edge.

Now suppose Λ contains a $W(\beta, \delta)$ with two β -edges and one δ -edge.

Lemma 5.11. *Λ^* does not contain a boundary cycle with respect to ω .*

Proof. The proof is completely similar as that of Lemma 5.8. \square

Lemma 5.12. *There is no source or sink dual to a white face in Λ^* with respect to ω .*

Proof. The proof is exactly the same as that of Lemma 5.6 with α and β interchanged. \square

Lemma 5.13. *There is no source or sink dual to a black face in Λ^* with respect to ω .*

Proof. The proof is exactly the same as that of Lemma 5.7 with α and β interchanged. \square

In conclusion, by Lemmas 5.11, 5.12, and 5.13 and the equation $\sum_{vertices} I(v) + \sum_{faces} I(f) = 2$, we have proved Theorem 5.1 for the case that Λ contains a $W(\beta, \delta)$ trigon with two β -edges and one δ -edge.

For the case of a $W(\alpha, \gamma)$ trigon, we can apply the same argument by interchanging α and β , and by exchanging γ and δ .

Finally, this completes the proof of Theorem 5.1.

Theorem 3.15 says that G_1 contains a $B(\alpha, \beta)$ face and a $W(\lambda, \mu)$ bigon, where $\{\lambda, \mu\} = \{\alpha, \delta\}, \{\alpha, \gamma\}, \{\beta, \delta\}$ or $\{\beta, \gamma\}$. We may assume without loss of generality that G_1 contains a $B(\alpha, \beta)$ face and a $W(\alpha, \delta)$ bigon.

The next proposition will be used to prove Theorem 5.17.

Proposition 5.14. *G_1 contains either a black bigon with two positive edges or a trigon.*

Remark: There are two types of trigons in G_1 ; one is a trigon with all positive edges and the other is a trigon with one positive edge and two negative edges.

To prove Proposition 5.14, we need two lemmas.

Lemma 5.15. *There is no bigon in G_1 whose edges are negative with a $(\epsilon_N, \epsilon'_S)$ corner and a $(\epsilon_S, \epsilon'_N)$ corner as shown in Figure 26 (i).*




FIGURE 26. A black bigon with negative edges and with a $(\epsilon_N, \epsilon'_S)$ corner and a $(\epsilon_S, \epsilon'_N)$ corner.

Proof. Let f be such a bigon as in the hypothesis. We assume without loss of generality that f is black. Recall that \widehat{F}_2 is an incompressible torus containing the graph G_2 and H_B is the intersection of V_2 and a black side of M . ∂f appears on $\widehat{F}_2 \cup H_B$ as in Figure 26 (ii).

The two endpoints of the negative edges ϵ and ϵ' of f separate ∂v_1 and ∂v_2 into two arcs respectively. Let a (resp. b) be the arc on ∂v_1 (resp. ∂v_2) which contains the endpoints of α -, β -edges at v_1 (resp. γ -, δ -edges at v_2). Then $\epsilon \cup a$ and $\epsilon' \cup b$ are circles which cobound an annulus on \widehat{F}_2 , denoted by A . Also the two corners of f separate $\overline{H_B - (v_1 \cup v_2)}$ into two disks; let D be one which contains a and b . Then $A \cup D$ is an once punctured Klein bottle. Therefore $A \cup D \cup f$ is a Klein bottle, which is a contradiction. \square

Lemma 5.16. *No white $(\epsilon_N, \epsilon'_N)$ or $(\epsilon_S, \epsilon'_S)$ corners exist in G_1 .*

Proof. Since there is a $W(\alpha, \delta)$ bigon, we have a $W(\alpha, \delta)$ corner and a $W(\delta, \alpha)$ corner. Then it follows from Lemma 3.3 that a white $(\epsilon_N, \epsilon'_N)$ or $(\epsilon_S, \epsilon'_S)$ corner cannot exist. \square

Proof of Proposition 5.14. Suppose for contradiction that G_1 does not contain a black bigon with positive edges and does not contain a trigon. Observe that Lemmas 5.15, 5.16 imply that G_1 doesn't contain a white bigon with two negative edges. Let \overline{G} be a connected component of the reduced graph \overline{G}_1 of G_1 . Note that \overline{G} is a graph on the sphere \widehat{F}_1 .

Claim. *Every vertex in \overline{G} has valency at least 4.*

Proof. Since G_1 does not contain a black bigon with positive edges and does not contain a white bigon with negative edges, it is easy to see that no vertex in \overline{G} has valency 1 or 2. Suppose a vertex v in \overline{G} has valency 3. Then there are three types of the local configuration at v in G_1 . See Figure 27.




FIGURE 27. A vertex of valency 3 in \overline{G} .

For the first type, since there is no black bigon with positive edges, the edges of the black bigon in Figure 27 (i) are negative. Thus the adjacent white bigon has two negative edges, which is a contradiction.

For the second type, all three white bigons in Figure 27 (ii) are $W(\alpha, \delta)$ bigons since there is no white bigon with negative edges in G_1 . Therefore there are two edges in the same edge class at the same label at v , which is a contradiction to Lemma 3.4.

If we have the third type as in Figure 27 (iii), all black bigons have negative edges. By Lemma 5.15, all the corners of these black bigons are either $B(\epsilon_N, \epsilon'_N)$ corners or $B(\epsilon_S, \epsilon'_S)$ corners. This condition forces one of the three white corners at v to be either a white $(\epsilon_N, \epsilon'_N)$ corner or a white $(\epsilon_S, \epsilon'_S)$ corner, which is impossible by Lemma 5.16.

Therefore, every vertex in \overline{G} has valency at least 4. \square

Let V, E, F be the number of vertices, edges and faces respectively in \overline{G} . Since no trigons exist in G_1 , no trigons exist in \overline{G} . So every face except one in \overline{G} has at least 4 edges. Hence, $4F - 4 \leq 2E$. Combined with $V - E + F = 2$, we get $2E \leq 4V - 4$. On the other hand, by the Claim, $4V \leq 2E$. This is a contradiction. \square

For the purpose of the next theorem, let $D^2(p, q)$ be a Seifert fibered space over the disk with two exceptional fibers of orders p and q respectively. Recall that $M(r_2)$ is a toroidal Dehn filling.

Theorem 5.17. *$M(r_2)$ is either $D^2(2, q) \cup_T D^2(2, s)$, $D^2(2, 3) \cup_T D^2(r, s)$ or $D^2(2, q) \cup_T D^2(3, s)$ where T is an incompressible torus in $M(r_2)$. Furthermore, the fibers of the two Seifert fibered subspaces of $M(r_2)$ intersect exactly once on T .*

Proof. G_1 contains a $W(\alpha, \delta)$ bigon and a $B(\alpha, \beta)$ face. Then Theorem 3.2 and Lemma 3.12 imply that $M(r_2)$ is $D^2(2, q) \cup_T D^2(r, s)$ where T is \widehat{F}_2 and the two fibers are induced from an $\alpha \cup \delta$ curve and an $\alpha \cup \beta$ curve respectively which intersect exactly once on T .

On the other hand, Proposition 5.14 guarantees the existence of either a black bigon with two positive edges or a trigon in G_1 . If G_1 contains a black bigon with two positive edges, $M(r_2)$ is $D^2(2, q) \cup_T D^2(2, s)$, which is the first case.

If G_1 contains a trigon with three positive edges, which is a good face, then by Theorem 3.2 one of orders of the four exceptional fibers is 3. Then $M(r_2)$ is either $D^2(2, 3) \cup_T D^2(r, s)$ or $D^2(2, q) \cup_T D^2(3, s)$ depending on which side the trigon lies.

The only remaining case is when G_1 has a trigon with two negative edges and one positive edge. Let f be a trigon with one λ -edge and two negative edges where $\lambda \in \{\alpha, \beta, \gamma, \delta\}$ as in Figure 28 (i). Then Lemma 3.3 (if f is black) and Lemma 5.16 (if f is white) imply that the (ϵ, ϵ') corner of f must be either a $(\epsilon_N, \epsilon'_N)$ corner or a $(\epsilon_S, \epsilon'_S)$ corner.

Recall that \hat{F}_2 is an incompressible torus containing the graph G_2 . Let \hat{H} be the intersection of V_2 and the side of $M(r_2)$ where f lies. ∂f appears on $\hat{F}_2 \cup \partial \hat{H}$ as in Figure 28 (ii). Let A be an annulus on \hat{F}_2 containing all the edges of f . Let N be the regular neighborhood of $\hat{H} \cup A \cup f$ in the side where f lies. Then it is easy to see that the fundamental group of N is isomorphic to the group $\langle x, y \mid x^2 y^3 = 1 \rangle$. Therefore N is homeomorphic to $D^2(2, 3)$. This implies the second case of $M(r_2)$. \square




FIGURE 28. A trigon with two negative edges and one positive edge.


6. A LINK SURGERY DESCRIPTION OF THE TOROIDAL DEHN FILLING

In this section, we interpret the toroidal Dehn filling $M(r_2)$ as the manifold obtained by Dehn surgery on some link L in S^3 by using a black good face and a white good face.

G_1 contains a $B(\alpha, \beta)$ face and a $W(\alpha, \delta)$ bigon. We denote them by f_1 and f_2 respectively. Note that a $B(\alpha, \beta)$ face is good by Lemma 3.11.

Then we are exactly in the same situation as in [10, Section 5]. Thus we go through the same terminology and arguments as in [10, Section 5].

Let f be a good (λ, μ) face in Λ . Then by Lemma 4.2 in [10], there are vertices x, y of f in Λ , representing corners of ∂f , such that the

FIGURE 29. The external interval of f .

edges of ∂f on G_2 are, up to homeomorphism, as in Figure 29. The interval $[x, y]$ on vertex v_i , $i = 1, 2$, that contains no edges of ∂f in its interior, is called *the external interval of f on vertex i* , denoted by I_f^i . In particular, if f is a bigon, there are two such intervals. Then we denote the external interval of f_i at vertex v_j by I_i^j , $j = 1, 2$. There are two choices on I_2^j :

- Choice (I). I_2^j doesn't contain the endpoints of the edge class β .
- Choice (II). I_2^j contains the endpoints of the edge class β .


Note that I_1^j contains the endpoints of the edge class δ . See Figure 30.

Recall that $M(r_2) = \widehat{M}_B \cup_{\widehat{F}_2} \widehat{M}_W$ and \widehat{H}_B (resp. \widehat{H}_W) = $V_2 \cap \widehat{M}_B$ (resp. \widehat{M}_W). For convenience, we change indices B, W into $1, 2$ respectively and we let V be the attached solid torus to M . Recall that r_1 and r_2 are slopes on $\partial_0 M = \partial V$ such that $M(r_1)$ is reducible and $M(r_2)$ is toroidal and the boundary components of F_1 have slopes r_1 .

Now follow the argument from the page 452 through the page 457 in [10, Section 5] with some notations changed, i.e. with $\epsilon, \delta_1, \delta_2, H_1, H_2, \mu, \tau, Q, T$ and M_i replaced by $\alpha, \beta, \delta, \widehat{H}_1, \widehat{H}_2, r_1, r_2, F_1, F_2$, and \widehat{M}_i .



Then from the page 456 and the page 457 in [10, Section 5], we have the following: *($M(r_2), K_{r_2}$) is obtained from (S^3, K_0) by some Dehn surgery on L_0 where K_{r_2} is the core of V in $M(r_2)$. That is, the exterior of K_{r_2} is obtained from the exterior of K_0 in S^3 by a Dehn surgery on L_0 . Moreover, given the 0-framing of K_0 in S^3 , the meridian of K_0 corresponds to the slope r_2 on ∂V .*

Here, as in Section 5 in [10], K_0 means the core of $V = \widehat{H}_1 \cup \widehat{H}_2 \subset S^3$ and L_0 means the link $K_1 \cup K'_1 \cup K_2 \cup K'_2$ in S^3 depicted in Figure 31. Also let L be the link $L = K_1 \cup K'_1 \cup K_2 \cup K'_2 \cup K_0$. Then L looks like Figure 32 according to the two choices (I), (II) on I_2^j . Note that L in Choice (II) is the reflection of L in Choice (I) (in which $K_i \mapsto K_i$, $0 \leq i \leq 2$, and $K'_i \mapsto K'_i$, $i = 1, 2$). Therefore under this reflection the

FIGURE 30. The external intervals of f_1 and f_2 .


r_1 -framing on K_0 goes to the $(-r_1)$ -framing (and the longitude to the longitude).

Let $L(\theta, \varphi, \psi, \omega, \pi)$ be the Dehn filling on the exterior of L where $\theta, \varphi, \psi, \omega, \pi$ denote the filling slopes on $K_1, K'_1, K_2, K'_2, K_0$ respectively. When one of the filling slopes is an asterisk (*), then no filling is done on the corresponding component.

FIGURE 31. The link $L_0 = K_1 \cup K'_1 \cup K_2 \cup K'_2$.FIGURE 32. The link $L = K_1 \cup K'_1 \cup K_2 \cup K'_2 \cup K_0$.

Proposition 6.1. $M = L(\theta, \varphi, \psi, \omega, *)$ where L is the link from Choice (I) as in Figure 32. Furthermore, the reducible Dehn filling $M(r_1)$ is $L(\theta, \varphi, \psi, \omega, 1/3)$ and the toroidal Dehn filling $M(r_2)$ is $L(\theta, \varphi, \psi, \omega, 1/0)$.

Proof. From the above discussion, we need only to show that $\pi = 1/3$ for $M(r_1)$. In other words, with respect to the framing of K_0 , the slope r_1 is $1/3$. Let a be a vertex in G_1 which contains a $B(\beta, \alpha)$ corner of a $B(\alpha, \beta)$ face f_1 . That is, v_1 in G_2 has the label a at the endpoint of a β -edge and v_2 in G_2 has the label a at the endpoint of an α -edge. Let

FIGURE 33. The r_1 -framing.

b be a vertex in G_1 which contains a $W(\alpha, \delta)$ corner of a $W(\alpha, \delta)$ bigon f_2 . Then it suffices to show that ∂b has slope $1/3$.

∂b consists of three black corners and three white corners, one of which is a $W(\alpha, \delta)$ corner. There are three occurrences of the label b at v_1 and v_2 . Let b_1, b_2 and b_3 be the labels representing three occurrences of b at v_1 and v_2 such that b_i at v_1 and b_i at v_2 are connected by a black corner of the vertex b , and b_i at v_2 and b_{i+1} at v_1 are connected by a white corner of the vertex b , $i = 1, 2, 3 \pmod{4}$. We assume that a $W(\alpha, \delta)$ corner connects b_3 at v_2 and b_1 at v_1 . The labels b_1, b_2 and b_3 on ∂v_1 and ∂v_2 occur either in the order $b_1 b_2 b_3$ or in the order $b_1 b_3 b_2$.

First, we assume the order $b_1 b_2 b_3$. Then we take Choice (I) for I_2^j . Since the r_1 -framings on \widehat{H}_1 and \widehat{H}_2 are shown Figure 5.16 and Figure 5.17 Case (I) in [10] respectively, a $B(\beta, \alpha)$ corner and a $W(\alpha, \delta)$ corner, which are uniquely determined on $\partial \widehat{H}_1$ and $\partial \widehat{H}_2$ respectively are given in Figure 33 (i). Therefore, ∂v has slope $1/3$.

Second, we assume the order $b_1 b_3 b_2$. Then we take Choice (II) for I_2^j . The r_1 -framing on \widehat{H}_2 is shown in Figure 5.17 Case (II) in [10]. Thus, ∂v has slope $-1/3$. See Figure 33 (ii). Under the reflection sending L in Choice (II) to L in Choice (I), $-1/3$ in Choice (II) goes to $1/3$ in Choice (I). Therefore we may assume without loss of generality that $\pi = 1/3$ in Choice (I). This completes the proof of $\pi = 1/3$. \square

Lemma 6.2. ψ is equal to 2 in Choice (I) and -2 in Choice (II).

Proof. ψ is the filling slope on K_2 , which is a core of V_2 disjoint from $\widehat{H}_2 \cup R_2$. Also $(W_2; \widehat{H}_2, R_2)$ is obtained from $(V_2; \widehat{H}_2, R_2)$ by ψ -Dehn filling on K_2 as described in the page 456 in [10]. Recall that $W_2 = \text{nbhd}(A_2 \cup \widehat{H}_2 \cup f_2)$. Since f_2 is a $W(\alpha, \delta)$ bigon, W_2 is the neighborhood of a Möbius band. Then with the standard framing on K_2 which is shown in Figure 5.17 Case (I) in [10], we easily see that $\psi = 2$ in Choice (I). For Choice (II), notice that the longitude has the opposite orientation to Choice (I). Therefore, $\psi = -2$ in Choice (II). \square

7. DEHN FILLINGS ON A LINK AND TANGLE FILLINGS ON A TANGLE

From now on, L is the link in Choice (I) as in Figure 32. Proposition 6.1 says that M , the reducible Dehn filling $M(r_1)$ and the toroidal Dehn filling $M(r_2)$ are obtained by Dehn surgery on L . L is strongly invertible and the quotient of the exterior under this involution gives rise to a tangle. Therefore M , $M(r_1)$ and $M(r_2)$ are double branched cover of S^3 along some tangles or some links which are obtained by rational tangle fillings on the tangle obtained from the quotient of the exterior of L .




FIGURE 34. The strong invertibility of L and the quotient of (S^3, L) .

We will use the same convention on rational tangles and the same terminology in [10, Section 6]. Let $T(\alpha_1, \dots, \alpha_n)$ be a filling of a tangle T where $\alpha_i \in \mathbf{Q} \cup \{1/0\}$, $1 \leq i \leq n$. Then the double branched cover of S^3 along the link $T(\alpha_1, \dots, \alpha_n)$ is a closed 3-manifold denoted by $\widehat{T}(\alpha_1, \dots, \alpha_n)$.

By Proposition 6.1, $M = L(\theta, \varphi, \psi, \omega, *)$. Figure 34 (i) shows that L is strongly invertible and depicts the quotient along with its fixed set. Then the quotient of (S^3, L) under this strong inversion is $P(\frac{0}{1}, \frac{1}{1}, \frac{0}{1}, \frac{1}{1}, \frac{0}{1})$ as in Figure 34 (ii), where P is the pentangle (See [10, Section 6]) for

the definition of the pentangle.) and invariant regular neighborhoods of L become tangle balls in the quotient. In other words, the quotient of the exterior $E(L)$ of L in S^3 under the involution is the pentangle P . The longitude framings of $\partial E(L)$ pass to the $\frac{1}{0}$ framings of the tangle balls as indicated by the dashed line. The meridian framings of $\partial E(L)$ pass to the $\frac{0}{1}$ framings of the tangle balls for K_0, K_1, K_2 and $\frac{1}{1}$ framings for K'_1, K'_2 as indicated by the solid line. In particular the (meridian, longitude)-framings of K_0, K_2 pass to the $(\frac{0}{1}, \frac{1}{0})$ -framing of the corresponding tangle spheres of P , which implies with Proposition 6.1 and Lemma 6.2 that the reducible Dehn filling $M(r_1) = L(\theta, \varphi, 2, \omega, 1/3)$ is $\tilde{P}(\theta', \varphi', 1/2, \omega', 3)$ and the toroidal Dehn filling $M(r_2) = L(\theta, \varphi, 2, \omega, 1/0)$ is $\tilde{P}(\theta', \varphi', 1/2, \omega', 0/1)$.




FIGURE 35. The pentangle $P(\theta', \varphi', 1/2, \omega', *) =$ the tangle $Q(-1/\theta', -1/\varphi', -1/\omega', *)$.

However we can regard the pentangle $P(\theta', \varphi', 1/2, \omega', *)$ as the tangle $Q(-1/\theta', -1/\varphi', -1/\omega', *)$ by rotating the pentangle by $-\pi/2$ where Q is the tangle as shown in Figure 35 (ii). Note that the tangle Q admits a symmetry by rotation in the horizontal axis. Without ambiguity, if we let θ, φ, ω be $-1/\theta', -1/\varphi', -1/\omega'$ respectively, then since M is $\tilde{P}(\theta', \varphi', 1/2, \omega', *)$, $M = \tilde{Q}(\theta, \varphi, \omega, *)$, $M(r_1) = \tilde{Q}(\theta, \varphi, \omega, -1/3)$ and $M(r_2) = \tilde{Q}(\theta, \varphi, \omega, 1/0)$.

Theorem 7.1. *M is the double branched cover of one of the tangles $Q(2, p-2, 1/p, *)$ where p is an integer and $p \leq -3$.*

To prove this, we do a series of computations describing the double branched covers of certain tangle filling of Q and combining this with certain Dehn filling theorems, we put restrictions on the tangle slopes θ, φ and ω .

Lemma 7.2. *Let M be an irreducible 3-manifold with ∂M a torus. Let θ, φ be slopes on ∂M such that $M(\theta)$ is reducible and $M(\varphi)$ is either reducible, a lens space, or S^3 . Then $\Delta(\theta, \varphi) \leq 1$.*

Proof. This is exactly Lemma 8.1 of [10]. □




FIGURE 36. $W(*) = Q(*, \varphi, \omega, 1/0)$ and the rational tangle fillings.

Lemma 7.3. $\theta, \varphi \notin \{-1, 0, 1, 1/0\}$ and $\omega \notin \{-1, 0, 1, 1/0, 1/2\}$.

Proof. Let $W(*)$ be $Q(*, \varphi, \omega, 1/0)$. Then $W(\theta) = Q(\theta, \varphi, \omega, 1/0)$ and $\widetilde{W}(\theta) = \widetilde{Q}(\theta, \varphi, \omega, 1/0) (= M(r_2))$. Theorem 5.17 says that $M(r_2) = \widetilde{W}(\theta)$ is a toroidal manifold and is not a Seifert fibered space. Moreover, by Lemma 7.2, $\widetilde{W}(\theta)$ is also irreducible since $\Delta(r_1, r_2) = \Delta(1/0, -1/3) = 3$. Consider the rational tangle fillings $W(-1)$, $W(0)$, $W(1)$ and $W(1/0)$. See Figure 36. Then the double branched cover of each tangle filling is either a Seifert fibered space or a reducible manifold, which is impossible. Therefore, $\theta \notin \{-1, 0, 1, 1/0\}$. On the other hand, the tangle Q admits a symmetry by rotation in the horizontal axis, which interchanges θ, φ . Therefore $\varphi \notin \{-1, 0, 1, 1/0\}$.

For ω , let $W'(*)$ be $Q(\theta, \varphi, *, 1/0)$. Then $M(r_2) = \widetilde{W}'(\omega)$ is a toroidal and irreducible manifold and is not a Seifert fibered space. Consider the




FIGURE 37. $W'(*) = Q(\theta, \varphi, *, 1/0)$ and the rational tangle fillings.


rational tangle fillings illustrated in Figure 37. The double branched cover of each tangle filling is either a Seifert fibered space or a reducible manifold, which is impossible. Hence $\omega \notin \{-1, 0, 1, 1/0\}$.

If $\omega = 1/2$, then $M(r_2) = \widetilde{W}'(1/2)$ contains a Klein bottle from figure 37, which is a contradiction. Thus $\omega \neq 1/2$. \square

Recall that $D^2(a, b)$ is a Seifert fibered space over D^2 with two exceptional fibers of orders a, b . Similarly, let $S^2(a, b, c)$ be a Seifert fibered space over S^2 with three exceptional fibers of orders a, b, c .

Lemma 7.4. *θ, φ are integers and ω is $1/k$ where k is an integer.*

Proof. First, we show that θ is an integer. Let $W(*) = Q(*, \varphi, \omega, -1/3)$. Then $\widetilde{W}(\theta) = \widetilde{Q}(\theta, \varphi, \omega, -1/3)$ is $M(r_1)$, which is reducible. We perform the rational tangle fillings shown in Figure 38. Observe that $\widetilde{W}(1/0)$ is either S^3 , $S^2 \times S^1$ or a lens space. Hence $\Delta(\theta, 1/0) \leq 1$ by Lemma 7.2, provided that $\widetilde{W}(*)$ is irreducible. Since $\theta \neq 1/0$ by Lemma 7.3, θ is an integer. Thus, to show that θ is an integer, it remains to show that $\widetilde{W}(*)$ is irreducible.



$$W(*) = Q(*, \varphi, \omega, -1/3)$$

FIGURE 38. $W(*) = Q(*, \varphi, \omega, -1/3)$ and the rational tangle fillings.

Suppose for contradiction that $\widetilde{W}(*)$ is reducible, i.e. $\widetilde{W}(*) = X \# Y(*)$ where Y contains the torus boundary component where Dehn fillings are performed. Note that if there are two slopes p and q such that $\widetilde{W}(p)$ and $\widetilde{W}(q)$ are prime manifolds, then $\widetilde{W}(p) = \widetilde{W}(q)$. However $\widetilde{W}(1/0)$ is either S^3 , $S^2 \times S^1$ or a lens space, which are all prime. Thus $\widetilde{W}(p)$ is either S^3 , $S^2 \times S^1$ or a lens space if $\widetilde{W}(p)$ is prime. From Figure 38, $\widetilde{W}(0)$ and $\widetilde{W}(-1)$ are prime manifolds because $\varphi, \omega \notin \{-1, 0, 1, 1/0\}$, whence they must be either S^3 , $S^2 \times S^1$ or a lens space.

Suppose $\varphi \neq 1 + 1/a$ for an integer a . If $\omega \neq 1/(1 + 1/b)$ for an integer b , then $\widetilde{W}(-1)$ is a toroidal prime manifold, which is neither S^3 , $S^2 \times S^1$ nor a lens space. If $\omega = 1/(1 + 1/b)$ for some integer b , then $\widetilde{W}(-1)$ is $S^2(e, f, g)$ where $e, f, g \geq 2$ since $\omega \neq 0, 1/2$, which is neither S^3 , $S^2 \times S^1$ nor a lens space, a contradiction. Therefore φ must be $1 + 1/a$ for some integer a . Then $\widetilde{W}(1) = S^2(|2a + 1|, 2, t)$ for some integer t . Note that $|2a + 1| \neq 0, 1$ because $\varphi \notin \{-1, 0, 1, 1/0\}$. If $\omega \neq 1/3$ i.e. $t \neq 0$, then $\widetilde{W}(1)$ is a prime manifold, which must be either S^3 , $S^2 \times S^1$ or a lens space. Hence ω must be $1/(3 + 1/s)$ for some integer s . This implies that $\widetilde{W}(-1) = S^2(|2a + 1|, 2, |2s + 1|)$. However,




FIGURE 39. $W'(*) = Q(\theta, \varphi, *, -1/3)$ and the rational tangle fillings.

since $\varphi \neq -1, 0, 1$ and $\omega \neq -1, 0, 1, 1/2$, $|2a+1| \geq 2$ and $|2s+1| \geq 2$, whence $\widetilde{W}(-1)$ is neither S^3 , $S^2 \times S^1$ nor a lens space, a contradiction. If $\omega = 1/3$, then $\widetilde{W}(-1) = S^2(|2a+1|, 2, 2)$, which is neither S^3 , $S^2 \times S^1$ nor a lens space, a contradiction. This completes the proof that $\widetilde{W}(*)$ is irreducible.

By the symmetry by rotation in the horizontal axis on the tangle Q , φ is also an integer.

To show that ω is $1/k$ for some integer k , a similar argument to the above applies. Let $W'(*) = Q(\theta, \varphi, *, -1/3)$. Then $\widetilde{W}'(\omega) = \widetilde{Q}(\theta, \varphi, \omega, -1/3)$ is $M(r_1)$, which is reducible. We perform the rational tangle fillings shown in Figure 39. Since $\widetilde{W}'(0)$ is either S^3 , $S^2 \times S^1$ or a lens space, ω is $1/k$ for some integer k by Lemma 7.2, provided that $\widetilde{W}'(*)$ is irreducible. However, $\widetilde{W}'(1/2)$ is prime since $\theta, \varphi \notin \{-1, 0, 1, 1/0\}$, and $\widetilde{W}'(1/2) \neq \widetilde{W}'(0)$. Thus, $\widetilde{W}'(*)$ is irreducible. The proof is complete. \square

Proposition 7.5. θ is 2 up to the symmetry by rotation in the horizontal axis on Q .

Before proving Proposition 7.5, we need to consider several special cases.

Lemma 7.6. $\theta, \varphi \neq -2$.

Proof. Suppose $\varphi = -2$. Let $W(*) = Q(*, -2, \omega, -1/3)$. Then $\widetilde{W}(\theta)$ is reducible. Consider the rational tangle fillings shown in Figure 38 with the -2 -rational tangle substituted for φ . Note that $\widetilde{W}(1)$ is a lens space or a reducible manifold. Since $\widetilde{W}(1/0)$ and $\widetilde{W}(-1)$ are prime and they are not homeomorphic, $\widetilde{W}(*)$ is irreducible. Therefore we can apply Lemma 7.2 to get $\Delta(\theta, 1) \leq 1$. Thus $\theta = 0, 1, 2$. However 0 and 1 are impossible by Lemma 7.3. Then $\theta = 2$. We will show that 2 is also impossible.

Consider the toroidal Dehn filling $M(r_2) = \widetilde{Q}(\theta, \varphi, \omega, 1/0)$. Since $\theta = 2$ and $\varphi = -2$, $M(r_2)$ contains a Klein bottle, which is a contradiction to the assumption that $M(r_2)$ does not contain a Klein bottle.

By the symmetry on Q , $\theta \neq -2$. \square

Lemma 7.7. If $\varphi = -3$, then $\theta = 2$.

Proof. Assume that $\varphi = -3$. Let $W(*) = Q(*, -3, \omega, -1/3)$. Then $\widetilde{W}(\theta)$ is reducible. Consider the rational tangle fillings shown in Figure 38 with the -3 -rational tangle substituted for φ .

Assume $\widetilde{W}(*)$ is hyperbolic. Notice that $\widetilde{W}(1) = S^2(2, 2, |1/\omega - 3|)$ and $\widetilde{W}(-1)$ is toroidal since $\omega \notin \{-1, 0, 1, 1/0, 1/2\}$. Then $\widetilde{W}(*)$ admits a reducible Dehn filling $\widetilde{W}(\theta)$, a toroidal Dehn filling $\widetilde{W}(-1)$, and a Dehn filling of the form $S^2(2, 2, 1/\omega - 3)$. By Corollary 1.3 in [12], $\Delta(\theta, 1) \leq 2$ and by Theorem 1.1 in [15], $\Delta(\theta, -1) \leq 3$. Thus $\theta = -1, 0, 1$ or 2 . However, Lemma 7.3 leads to $\theta = 2$. The proof is complete under the assumption that $\widetilde{W}(*)$ is hyperbolic.

We show that $\widetilde{W}(*)$ is hyperbolic. Equivalently, it suffices to show that $\widetilde{W}(*)$ is irreducible, ∂ -irreducible, non-Seifert fibered and atoroidal. Since $\widetilde{W}(1/0)$ and $\widetilde{W}(-1)$ are prime and are not homeomorphic, $\widetilde{W}(*)$ is irreducible.

Suppose $\widetilde{W}(*)$ is a Seifert fibered space. Since $\widetilde{W}(1/0)$ is a lens space, $\widetilde{W}(*)$ is a Seifert fibered space over D^2 with at most 2 exceptional fibers. Hence $\widetilde{W}(r)$ is a Seifert fibered space over S^2 with at most three exceptional fibers except for one slope r , for which $\widetilde{W}(r)$ is reducible. Figure 38 shows that $\widetilde{W}(-1)$ is irreducible. Hence $\widetilde{W}(-1) = S^2(e, f, g)$ where $e, f, g \geq 1$, which doesn't contain a separating incompressible torus. On the other hand, $\widetilde{W}(-1)$ contains a separating incompressible torus which is the double branched cover of S in Figure 38, since $\omega \notin$

$\{-1, 0, 1, 1/0, 1/2\}$. This is a contradiction. Therefore $\widetilde{W}(\ast)$ is not a Seifert fibered space.

Suppose $\widetilde{W}(\ast)$ is ∂ -irreducible. After ∂ -compression, the torus boundary becomes a sphere which must bound a 3-ball since $\widetilde{W}(\ast)$ is irreducible. This implies that $\widetilde{W}(\ast)$ is a solid torus, which is a Seifert fibered space, a contradiction.

Suppose $\widetilde{W}(\ast)$ is toroidal. Let T be an essential torus in $\widetilde{W}(\ast)$. Since $\widetilde{W}(1/0)$ is a lens space, T must be separating, which allows us to let $\widetilde{W}(\ast) = A \cup_T B(\ast)$ where B contains the torus boundary of $\widetilde{W}(\ast)$.

Claim. T is compressible in $\widetilde{W}(-1)$.

Proof. Suppose T is incompressible in $\widetilde{W}(-1)$. Figure 38 shows that there is a unique incompressible torus up to isotopy in $\widetilde{W}(-1)$, which must be the double branched cover of the sphere S as in Figure 38. Therefore $\widetilde{W}(-1) = D^2(2, 4) \cup_T D^2(2, |1/\omega - 1|)$. Thus A is either $D^2(2, 4)$ or $D^2(2, |1/\omega - 1|)$. Observe from Figure 38 that $\widetilde{W}(0) = S^2(3, 3, |1/\omega - 2|)$, $\widetilde{W}(1) = S^2(2, 2, |1/\omega - 3|)$ and $\widetilde{W}(2) = S^2(3, 2, |1/\omega + 1|)$. Hence A is $D^2(2, 3)$ and $\omega = 1/4$. $A = D^2(2, 3)$ is the complement of the trefoil knot i.e. the $(2, 3)$ torus knot $T_{2,3}$. $\widetilde{W}(1) = S^2(2, 2, 1)$ becomes a lens space. It is easy to see that $\widetilde{W}(1) = L(8, -3)$. Since $B(1)$ is a solid torus, $\widetilde{W}(1) = L(8, -3)$ is obtained by Dehn filling on $A = T_{2,3}$. This is impossible by Corollary 7.4 in [4]. \square

T is compressible in $\widetilde{W}(1/0)$, $\widetilde{W}(0)$, $\widetilde{W}(1)$, $\widetilde{W}(-1)$ and $\widetilde{W}(2)$. Since $\Delta(2, -1) = \Delta(2, 0) = \Delta(1, -1) \geq 2$, by Lemma 2.4 in [2] $B(\ast)$ is a cable space $C(p, q)$ with the cabling slope $1/0$. (See Section 3 in [5] for the definition of a cable space.). Moreover, since $\Delta(\theta, 1/0) = 1$, $B(\theta)$ is a solid torus, which makes T compressible in $\widetilde{W}(\theta)$. Therefore $\widetilde{W}(1/0) = A(r_{1/0}) \# L(q, p)$ and $\widetilde{W}(\theta) = A(r_\theta)$ for some slopes $r_{1/0}, r_\theta$ on T with $\Delta(r_{1/0}, r_\theta) = |q|\Delta(\theta, 1/0) = |q|$. Since $\widetilde{W}(1/0)$ is a lens space, $A(r_{1/0})$ must be S^3 . Then $|q|$ must be 1 by Lemma 7.2 since A is irreducible, $A(r_{1/0})$ is S^3 and $A(r_\theta)$ is reducible. The fact that $q = 1$ or -1 implies that T is ∂ -parallel in $\widetilde{W}(\ast)$, which is a contradiction to the assumption that T is essential in $\widetilde{W}(\ast)$. Thus $\widetilde{W}(\ast)$ is atoroidal. \square

Lemma 7.8. *If $\varphi = 3$, then $\theta = 2$.*

Proof. Assume that $\varphi = 3$. Let $W(\ast) = Q(\ast, 3, \omega, -1/3)$. Then $\widetilde{W}(\theta)$ is reducible. Consider the rational tangle fillings shown in Figure 38 with the 3-rational tangle substituted for φ .

Assume $\widetilde{W}(\ast)$ is hyperbolic. $\widetilde{W}(-1)$ is toroidal since $\omega \notin \{-1, 0, 1, 1/0, 1/2\}$. $\widetilde{W}(\theta)$ is reducible. Thus $\Delta(\theta, -1) \leq 3$. The possible values of θ are $-4, -3, -2, 2$ since $\theta \notin \{-1, 0, 1, 1/0\}$. If θ is 2, we are done. If θ is -3 , then this belongs to the case of $\varphi = -3$ by the symmetry on Q , whence we are done. Also, $\theta \neq -2$ by Lemma 7.6. The remaining case is $\theta = -4$.

Suppose $\theta = -4$. Let $W'(\ast) = Q(-4, 3, \ast, -1/3)$. Then $\widetilde{W}'(\omega)$ is reducible. Similarly, we can easily see from Figure 39 that $\widetilde{W}'(1/0)$ is the lens space $L(7, -2)$ and $\widetilde{W}'(1)$ is $S^2(4, 2, 3)$. They are non-homeomorphic prime manifolds. Thus $\widetilde{W}'(\ast)$ is irreducible. Moreover, Lemma 7.2 leads to $\Delta(1/0, \omega) \leq 1$. Since ω is of the form $1/k$ for some integer k , ω is 1 or -1 , which is impossible by Lemma 7.3. Hence $\theta \neq -4$.

To complete the proof of the lemma, it remains to show that $\widetilde{W}(\ast)$ is hyperbolic.

$\widetilde{W}(\ast)$ is irreducible because $\widetilde{W}(1/0)$ and $\widetilde{W}(-1)$ are prime and are not homeomorphic. The way of proving that $\widetilde{W}(\ast)$ is ∂ -irreducible and not a Seifert fibered space is exactly the same as for the case $\varphi = -3$.

Suppose $\widetilde{W}(\ast)$ is toroidal. Let T be an essential torus in $\widetilde{W}(\ast)$. Since $\widetilde{W}(1/0)$ is a lens space, T must be separating. Let $\widetilde{W}(\ast) = A \cup_T B(\ast)$ where B contains the torus boundary of $\widetilde{W}(\ast)$.

Claim. T is compressible in $\widetilde{W}(-1)$.

Proof. Suppose T is incompressible in $\widetilde{W}(-1)$. Let T' be the double branched cover of S as shown in Figure 38. $\widetilde{W}(-1) = D^2(2, 2) \cup_{T'} D^2(2, |1/\omega - 1|)$. Then T' is the unique incompressible torus up to isotopy in $\widetilde{W}(-1)$ unless $\omega \in \{-1, 0, 1, 1/0, 1/2, 1/3\}$. If $\omega = 1/3$, then $\widetilde{W}(1)$ is reducible from Figure 38. Thus by Lemma 7.2, $\Delta(\theta, 1) \leq 1$ i.e. $\theta = 2$ since $\theta \neq 0, 1$, in which case we are done. Therefore we may assume that T' is the unique incompressible torus up to isotopy in $\widetilde{W}(-1)$. Thus, $T = T'$. In other words, $\widetilde{W}(-1) = D^2(2, 2) \cup_T D^2(2, |1/\omega - 1|)$. Hence A is either $D^2(2, 2)$ or $D^2(2, |1/\omega - 1|)$. Observe from Figure 38 that $\widetilde{W}(0) = S^2(3, 3, |1/\omega - 2|)$, $\widetilde{W}(1) = S^2(4, 2, |1/\omega - 3|)$ and $\widetilde{W}(2) = S^2(3, 2, |1/\omega - 5|)$. Hence A is $D^2(2, 3)$ and $\omega = 1/4$. $A = D^2(2, 3)$ is the complement of the trefoil knot i.e. the $(2, 3)$ torus knot $T_{2,3}$. Now apply the same argument as in the case $\varphi = -3$ using $\widetilde{W}(1) = L(10, -3)$. \square

T is compressible in $\widetilde{W}(1/0)$, $\widetilde{W}(0)$, $\widetilde{W}(1)$, $\widetilde{W}(-1)$ and $\widetilde{W}(2)$. Then the rest of the proof is exactly the same as for the case $\varphi = -3$. \square

Lemma 7.9. *If $\omega = -1/3$, then $\theta = 2$.*

Proof. Assume that $\omega = -1/3$. Let $W(*) = Q(\theta, *, -1/3, -1/3)$. Then $\widetilde{W}(\varphi)$ is reducible. Consider the rational tangle fillings shown in Figure 40 with the $-1/3$ -rational tangle substituted for ω . Then $\widetilde{W}(-1)$ is toroidal unless $\theta = 1/0, 0, 1, 2$ (by Lemma 7.3, $\theta = 2$, in which case we are done.). Hence we may assume that $\widetilde{W}(-1)$ is toroidal.

Assume $\widetilde{W}(*)$ is hyperbolic. $\widetilde{W}(\varphi)$ is reducible and $\widetilde{W}(-1)$ is toroidal. Thus $\Delta(\varphi, -1) \leq 3$. By Lemma 7.3 $\varphi = -4, -3, -2$ or 2 . If $\varphi = -3$, then $\theta = 2$ by Lemma 7.7. If $\varphi = 2$, we are done by using the symmetry by rotation in the horizontal axis on Q . Also, $\varphi \neq -2$ by Lemma 7.6.

Suppose $\varphi = -4$. Let $W'(*) = Q(*, -4, -1/3, -1/3)$. Then $\widetilde{W}'(\theta)$ is reducible. Similarly, we can easily see from Figure 38 with W replaced by W' that $\widetilde{W}'(2)$ is the lens space $L(5, 1)$ and $\widetilde{W}'(0)$ is $S^2(3, 4, 5)$. These are non-homeomorphic prime manifolds. Thus $\widetilde{W}'(*)$ is irreducible. Since $\widetilde{W}'(2)$ is a lens space and $\widetilde{W}'(\theta)$ is reducible, by Lemma 7.2, $\Delta(\theta, 2) \leq 1$. Thus θ is $1, 2$ or 3 . However 1 is ruled out by Lemma 7.3. If $\theta = 2$, then we are done. The case $\theta = 3$ belongs to the case $\varphi = 3$ by the symmetry on Q . Thus $\theta = 2$ by Lemma 7.8.

To complete the proof of the lemma, it remains to show that $\widetilde{W}(*)$ is hyperbolic.

$\widetilde{W}(*)$ is irreducible because $\widetilde{W}(1/0)$ and $\widetilde{W}(-1)$ are prime and not homeomorphic. In the proof of Lemma 7.7, in order to prove that $\widetilde{W}(*)$ is not a Seifert fibered space, we used the fact that $\widetilde{W}(-1)$ is irreducible and contains a separating incompressible torus. In this case, $\widetilde{W}(-1)$ is also irreducible and contains a separating incompressible torus unless $\theta = -1, 0, 1, 2$, in which cases we are done by Lemma 7.3. Now we can apply the same argument as in the case $\varphi = -3$ to show that $\widetilde{W}(*)$ is ∂ -irreducible and not a Seifert fibered space.

Suppose $\widetilde{W}(*)$ is toroidal. Let T be an essential torus in $\widetilde{W}(*)$. Since $\widetilde{W}(1/0)$ is a lens space, T must be separating. Let $\widetilde{W}(*) = A \cup_T B(*)$ where B contains the torus boundary of $\widetilde{W}(*)$.

Claim. *T is compressible in $\widetilde{W}(-1)$.*

Proof. Suppose T is incompressible in $\widetilde{W}(-1)$. Let T' be the double branched cover of S as shown in Figure 40. $\widetilde{W}(-1) = D^2(2, 4) \cup_{T'} D^2(2, |\theta - 1|)$. Then T' is the unique incompressible torus up to isotopy in $\widetilde{W}(-1)$ unless $\theta = -1, 0, 1, 1/0, 2$, in which cases we are done by Lemma 7.3. Therefore $T' = T$. That is, $\widetilde{W}(-1) = D^2(2, 4) \cup_T D^2(2, |\theta - 1|)$. Thus A is either $D^2(2, 4)$ or $D^2(2, |\theta - 1|)$. Observe from




FIGURE 40. $W(*) = Q(\theta, *, \omega, -1/3)$ and the rational tangle fillings.

Figure 40 that $\widetilde{W}(0) = S^2(3, 5, |\theta|)$, $\widetilde{W}(1) = S^2(2, 6, |\theta + 1|)$ and $\widetilde{W}(2) = S^2(3, 2, |\theta + 5|)$. Hence A is $D^2(2, 3)$ and $\theta = -2$. $A = D^2(2, 3)$ is the complement of the trefoil knot i.e. the $(2, 3)$ torus knot $T_{2,3}$. Now apply the same argument as in the case $\varphi = -3$ using $\widetilde{W}(1) = L(16, -5)$. \square

T is compressible in $\widetilde{W}(1/0)$, $\widetilde{W}(0)$, $\widetilde{W}(1)$, $\widetilde{W}(-1)$ and $\widetilde{W}(2)$. Then the rest of the proof is exactly the same as for the case $\varphi = -3$. \square

Lemma 7.10. $\omega \neq 1/3$.

Proof. Suppose that $\omega = 1/3$. Let $W(*) = Q(\theta, *, 1/3, -1/3)$. Then $\widetilde{W}(\varphi)$ is reducible. Consider the rational tangle fillings shown in Figure 40 with the $1/3$ -rational tangle substituted for ω . Note that $\widetilde{W}(0)$ is a lens space and $\widetilde{W}(1)$ is either a reducible manifold or a lens space. Therefore Lemma 7.2 says that $\Delta(\varphi, 0) \leq 1$ and $\Delta(\varphi, 1) \leq 1$ provided that $\widetilde{W}(*)$ is irreducible. Hence φ is 0 or 1 which is impossible by Lemma 7.3. However $\widetilde{W}(0)$ and $\widetilde{W}(-1)$ are prime manifolds and are not homeomorphic. This implies that $\widetilde{W}(*)$ is irreducible. \square

Proof of Proposition 7.5. Theorem 5.17 says that the toroidal manifold $M(r_2)$ is either $D^2(2, q) \cup_T D^2(2, s)$, $D^2(2, 3) \cup_T D^2(r, s)$ or $D^2(2, q) \cup_T D^2(3, s)$. On the other hand, $M(r_2)$ is the double branched cover $\tilde{Q}(\theta, \varphi, \omega, 1/0)$ over the link $Q(\theta, \varphi, \omega, 1/0)$ as shown in Figure 41. Let S be the sphere shown in Figure 41. The double branched cover of S is an incompressible torus in $M(r_2)$. Since there is the unique incompressible torus in $M(r_2)$ up to isotopy, T is the double branched cover of S . Therefore, by Lemma 7.4 $M(r_2) = D^2(|\theta|, |\varphi|) \cup_T D^2(2, |1/\omega|)$.




FIGURE 41. $Q(\theta, \varphi, \omega, 1/0)$.

(1) Assume $M(r_2) = D^2(2, q) \cup_T D^2(2, s)$. Then θ must be 2 or -2 . Lemma 7.6 implies that θ is 2.

(2) Assume $M(r_2) = D^2(2, q) \cup_T D^2(3, s)$. Then there are two cases: either $\{|\theta|, |\varphi|\} = \{3, s\}$ or $\{2, |1/\omega|\} = \{3, s\}$.

If $\{|\theta|, |\varphi|\} = \{3, s\}$, then up to the symmetry, we may assume that $|\varphi| = 3$. Therefore by Lemmas 7.7, 7.8, θ is 2.

If $\{2, |1/\omega|\} = \{3, s\}$, then $|1/\omega| = 3$. Lemmas 7.9, 7.10 imply that θ is 2.

(3) Assume $M(r_2) = D^2(2, 3) \cup_T D^2(r, s)$. There are two cases: $\{|\theta|, |\varphi|\} = \{2, 3\}$ or $\{2, |1/\omega|\} = \{2, 3\}$. In either case, Lemmas 7.7, 7.8, 7.9 and 7.10 show that θ is 2. \square

Proof of Theorem 7.1. Proposition 7.5 says that $\theta = 2$. Then $\tilde{Q}(2, \varphi, \omega, -1/3)$ is $S^2(2, 3, |\varphi + 2 - 1/\omega|)$. See $W(2)$ in Figure 38. However since $M(r_1) = \tilde{Q}(2, \varphi, \omega, -1/3)$ is reducible, $\varphi + 2 - 1/\omega$ must be 0. Thus if we let $\omega = 1/p$ where p is an integer, then $\varphi = p - 2$. Hence $M(r_2) = \tilde{Q}(2, p - 2, 1/p, -1/3)$. However Lemma 7.3 implies that $p \neq -1, 0, 1, 2, 3$. If p is either -2 or 4 , then $M(r_2) = \tilde{Q}(2, p - 2, 1/p, -1/3)$ contains a Klein bottle, which is a contradiction. Therefore $p \leq -3$ or $p \geq 5$. Observe that the tangle $Q(2, p - 2, 1/p, -1/3)$ is isotopic to

the tangle $Q(2, -p, 1/(-p+2), -1/3)$. Therefore, we may assume that $p \leq -3$.

To complete the proof of the theorem, we need only to show that $\tilde{Q}(2, p-2, 1/p, *)$ is hyperbolic. However, the proof of this is exactly the same as the proof of Theorem 4.2 in [2]. □

Acknowledgements

The author would like to thank Sangyop Lee for his useful conversations. Also, the author is very grateful to his advisor, Cameron McA Gordon, for his excellent suggestions and encouragements through this work. We should also mention that since the author finished writing this paper, he has learned that the main result of this paper has also recently been proved by Sangyop Lee independently.

REFERENCES

- [1] M. Culler, C. McA. Gordon, J. Luecke, and P.B. Shalen, *Dehn surgery on knots*, Ann. of Math. **125** (1987), 237–300.
- [2] M. Eudave-Muñoz and Y.Q. Wu, *Nonhyperbolic Dehn fillings on hyperbolic 3-manifolds*, Pacific J.Math. **190** (1999), 261–275.
- [3] C. McA. Gordon, *Boundary slopes on punctured tori in 3-manifolds*, Trans. Amer. Math. Soc. **350** (1998), 1713–1790.
- [4] C. McA. Gordon, *Dehn surgery and satellite knots*, Trans. Amer. Math. Soc. **275** (1983), 687–708.
- [5] C. McA. Gordon and R.A. Litherland, *Incompressible planar surfaces in 3-manifolds*, Topology Appl. **18** (1984), 121–144.
- [6] C. McA. Gordon and J. Luecke, *Knots are determined by their complements*, J. Amer. Math. Soc. **2** (1989), 371–415.
- [7] C. McA. Gordon and J. Luecke, *Dehn surgeries on knots creating essential tori, I*, Communications in Analysis and Geometry **3** (1995), 597–644.
- [8] C. McA. Gordon and J. Luecke, *Reducible manifolds and Dehn surgery*, Topology **35** (1996), 385–409.
- [9] C. McA. Gordon and J. Luecke, *Toroidal and boundary-reducing Dehn fillings*, Topology Appl. **93** (1999), 77–90.
- [10] C. McA. Gordon and J. Luecke, *Non-integral toroidal Dehn surgeries*, Communications in Analysis and Geometry **12** (2004), 417–485.
- [11] W. Jaco, *Lectures on three-manifold topology*, Regional Conf. Series in Math. **43**(1997).
- [12] S. Lee, *Reducing and Toroidal Dehn fillings on 3-manifold bounded by two tori*, Mathematical Research Letters **12**(2005) 10001–10014.
- [13] S. Lee, *Toroidal Dehn surgeries on Knots in $S^1 \times S^2$* , J. Knot Theory Ramifications **14**(2005) 657–664.
- [14] S. Lee, S. Oh and M. Teragaito, *Dehn fillings and small surfaces*, Preprint 2003. <http://de.arxiv.org/abs/math.GT/0303156>.
- [15] S. Oh, *Reducible and toroidal manifolds obtained by Dehn fillings*, Topology Appl. **75** (1997), 93–104.

- [16] Y.Q. Wu, *The reducibility of surgered 3-manifolds*, Topology Appl. **43** (1992), 213–218.
- [17] Y.Q. Wu, *Dehn fillings producing reducible manifolds and toroidal manifolds*, Topology **37** (1998), 95–108.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TEXAS AT AUSTIN, 1 UNIVERSITY STATION C1200 AUSTIN, TX 78712-0257

E-mail address: skang@math.utexas.edu